

BUTOXONE S. B.

For Use as a Foliage Spray on Soybeans to Control Cocklebur

ACTIVE INGREDIENT:

4-(2,4-Dichlorophenoxy) butyric acid,
dimethylamine salt*..... 26.3%
INERT INGREDIENTS..... 73.7%

*Equivalent to 22.2% by weight 4-(2,4-dichlorophenoxy) butyric acid.

U. S. Patent No. 2,863,754

GENERAL INFORMATION: Butoxone S.B. is specially recommended for use on soybeans wherever cocklebur (*Xanthium* spp.) is a serious weed problem. At the rates recommended for controlling cocklebur, certain other broadleaved weeds, such as pigweed (*Amaranthus* spp.) and annual morning glory (*Ipomoea* spp.) will also be suppressed, or partially controlled.

DIRECTIONS

WHEN TO USE: Make application from 7 to 10 days pre-bloom to mid-bloom period of the soybeans. Do not apply earlier than this stage of growth.

AMOUNT TO USE: Mix Butoxone S.B. at rate of 1 gal. to 120 gals. of water and apply 10 to 12 gals. of diluted spray per acre. It is important that sprayer be calibrated accurately before application.

HOW TO APPLY: Apply as foliage spray. Use ground sprayer with flat fan type nozzles. Use one nozzle over each row and adjust to a height above the crop to give overall coverage.

CAUTION: Avoid contact with skin, eyes or clothing. Do not use so as to contaminate water intended for irrigation or domestic use. Do not re-use shipping containers. Do not store near fertilizers, insecticides, fungicides or seeds. Store at temperatures above 32°F. If allowed to freeze, re-mix before using.

Avoid spray drift to susceptible plants and crops as it may injure cotton, tomatoes, ornamentals, etc. Do not use same spray equipment for other purposes without first thoroughly cleaning with alkali and water. Small amounts of 2,4-D, MCP, silvex or 2,4,5-T may damage soybeans; sprayers previously used to apply these chemicals must be thoroughly cleaned with alkali and water before being used for applying Butoxone S.B. Contact your supplier for special instructions on cleaning spray equipment.

Seller makes no warranty of any kind, express or implied, concerning the results of use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

Do NOT Store with Foodstuffs

Manufactured by

CHIPMAN
Bound Brook, N.J.
N.Kansas Ctiy, Mo.

CHEMICAL
Chicago, Ill.
Portland, Ore.

COMPANY,
Pasadena, Tex.
Bessemer, Ala.

INC.
Palo Alto, Calif.
St. Paul, Minn.

