



PRODUCT NO.  
**2600**

# RHO **2,4-D BUTYL** A SELECTIVE

**For control of broadleaf weeds and woody plants suscep**

**ACTIVE INGREDIENT:**

Butyl ester of 2,4-dichlorophenoxyacetic acid\* 49.0%

**INERT INGREDIENTS** 51.0%

Equivalent to 39.08% of 2,4-dichlorophenoxyacetic acid or not less than 3.34 lbs. of 2,4-dichlorophenoxyacetic acid per gal. at 68°F.

## **CAUTION: Keep Out of Reach of Children**

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, or clothing. In case of contact flush eyes with plenty of water; wash skin with soap and water. Avoid breathing spray mist.

Do not allow solution or spray drift to contact desirable plants. Among crops highly susceptible to 2,4-D injury are cotton, tobacco, blackeyed peas, beans, tomatoes, melons, other vegetables, grapes, fruit trees, and some ornamentals. (Coarse sprays are less likely to drift than are fine mist sprays.) It is difficult to completely clean 2,4-D from sprayers or other equipment, therefore do not use 2,4-D sprayers and other equipment for purposes where even trace amounts of this chemical may cause injury. Do not reuse shipping containers; destroy by crushing and burying. Avoid contamination of water intended for irrigation and domestic use. Do not transport with or store near seeds, fertilizers, insecticides, or fungicides. Vapors from this product may injure susceptible plants in the vicinity. Store at temperatures above 32°F. If allowed to freeze, roll drum to remix before using.

Do not use this product in vicinity of crop plants highly susceptible to 2,4-D injury. Do not use or store near heat or open flame.

Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers.

Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D.

Do not feed treated straw to livestock.

### **CONTROLS THESE WEEDS**

Dandelion	Pennyworts	Bull Thistle	Vetch	Tarweed or Fiddleneck
Plantains	Mustards	Poisonhemlock	Burdock	(spray when small)
Healall	Gumweed	Puncturevine	Ground Ivy	Ragweeds
Indian Mallow	Chicory	Yellow Starthistle	Pigweed	Stinging Nettle
or Velvetleaf	Wild Licorice	Wild Radish	Galinsoga	and many others

**NOTE:** These suggestions for use are general. Local conditions may affect the use of this chemical. Consult State Agricultural Extension authorities for specific recommendations as to dosage and time of application and for control of other weeds.

### **SELECTIVE WEEDING IN CROPS**

**ESTABLISHED PASTURE, FIELD, RANGE, AND TURF GRASSES:** Use 1 to 5 pts. per acre, depending on susceptibility of weeds. Apply in sufficient water for uniform coverage of weeds. Use low rate for easy-to-kill annuals and apply while weeds are small; repeat if new weeds appear. Spring or fall applications may be made, except on grass seed fields where fall applications are not recommended. Use 2 to 5 pts. per acre for perennial weeds such as Bindweed, Canada Thistle, St. Johnswort, and Garlic. In general spray these from vigorous spring growth to early bloom and again on fall regrowth. Repeat application for 2 or more successive years may be needed to

Do not graze dairy animals within 7 days after application

EPA Reg. No. 359-523  
EPA Est. 359-MO-1  
EPA Est. 359-OR-1

PL-1074

**ED**

Manuf  
**RHOD**  
**AGRICULTUR**  
SOMERSET,