

[359-526]
1 of 2

CHIPMAN MALATHION FLOWABLE 8

ACCEPTED
MAY 11 1971
FOR
ED
TO ATTACH

102
130 JEROME AVENUE, P.O. BOX 2000
NEW BRUNSWICK, NEW JERSEY 08903

RECEIVED

January 20, 1971

PRODUCT NO.

281

[359-526]
2 of 2

CHIPMAN[®] MALATHION FLOWABLE 8

NET CONTENTS
GALLONS

For use in controlling Spider Mites and insect pests of fruit, nut, vegetable crops and ornamentals. Also for controlling Chicken Mites and lice on poultry and in poultry houses.

ACTIVE INGREDIENT:

Malathion* 80.8%

INERT INGREDIENTS 19.2%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate, equivalent to 8 lb. per gallon at 68°F.

This formulation will not freeze at 0°C, or at -10°C. (14°F.), at -10° C. it will become somewhat viscous, but is not damaged by lower temperatures and will return to normal physical condition without agitation on standing at warmer temperature.

CAUTION: Keep Out of Reach of Children

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Destroy container when empty; never re-use. Punch container full of holes or crush it so it cannot be used or salvaged for any purpose.

To protect bees and pollinating insects, do not apply to crops in bloom nor to bee yards areas. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is poisonous to fish. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

PRECAUTIONS: Avoid contamination of feed and feeding troughs and water fountains with spray. Use a high pressure sprayer for surface sprays.

Local conditions may affect the use of this insecticide. Consult State agricultural extension or Experiment Station specialists for specific recommendations consistent with label for local pest problems.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Do NOT Store with Foodstuffs

DIRECTIONS: Apply when significant populations develop or according to local recommendations. Unless otherwise specified, mix with sufficient water for thorough, full-coverage applications.

| CROP | PESTS | AMOUNT | TIMING AND COMMENTS |
|---|---|------------------------------------|--|
| | Bud Moths, Woolly and Green Apple aphids, Rosy Apple Aphid, Forbes Scale | 10 oz. to 1 pt. per 100 gals. | Make at least two applications for Willamette and Two-Spotted Mites, 10 to 12 days apart in summer months. For Forbes Scale make two or three applications. Do not apply within 3 days of harvest. |
| APPLES | European Red, Clover, Willamette, Two-Spotted Mites, Mealybugs | 10 oz. to 1 1/4 pts. per 100 gals. | |
| | Codling Moth, Plum Curculio, Red Banded Leaf Roller | 1 1/4 pints per 100 gals. | |
| CHERRIES | Black Cherry Aphid, Cherry Fruit Flies, Bud Moth. | 10 oz. to 1 pt. per 100 gals. | For Cherry Fruit Flies, repeat application every 10 days, but not later than 3 days before harvest. |
| PEACHES | Oriental Fruit Moth, Plum Curculio | 1 1/4 pints per 100 gals. | Do not apply within 7 days of harvest. |
| | Mites, Pear Psylla, Mealybugs | 10 oz. to 1 1/4 pts. per 100 gals. | Do not apply within 1 day of harvest. Injury may occur to Bosc pears under certain conditions in the Northwest. |
| PEARS | Codling Moth, Plum Curculio, Fruit Tree and Red Banded Leafrollers | 1 1/4 pints per 100 gals. | |
| | Walnut Husk Flies | 2 to 3 pints | Apply in 500 gals. water per acre by air carrier type equipment. |
| | | 4 oz. to 6 oz. | Apply in 100 gals. by conventional high pressure spray equipment. |
| WALNUTS | Aphids and Mites | 1 to 2 pints | Apply in 400 gals. water per acre by air carrier type equipment. |
| | | 2 oz. to 4 oz. | Apply in 100 gals. by conventional high pressure spray equipment. |
| PEAS | Pea Aphid, Pea Leaf Weevil | 1 to 1 1/4 pts. 'A' | Use high rate for weevil. Make one or more applications as needed. Do not apply within 3 days of harvest. |
| STRAWBERRIES | Aphids, Spider Mites | 1 pt. per acre | Do not apply within 3 days of harvest. |
| CRANBERRIES | Leafhoppers, Black-Headed Fireworms, Spittlebug Nymphs, Cranberry Fruitworm | 1 pt. per acre | Do not apply within 3 days of harvest. |
| BRAMBLES (Blackberry, dew-berry, raspberry) | Aphids, Rose Scale | 1 3/4 pts. per acre | Do not apply within 3 days of harvest. |
| POTATOES | Aphids, Leafhoppers, Mealybugs | 10 oz. per acre | Resistant to mealybugs. |
| | False Chinch Bugs | 1 pt. per acre | |
| ORNAMENTALS | Aphids, Spider Mites | 1 1/4 pints per 100 gals. | Use only on certain ornamentals. Do not apply to Citrus, Ficus, and other plants. |
| POULTRY | Northern Fowl Mites, Poultry Lice | 5 1/2 oz. per 1 gal. water | Use 1 gal. mixture per 100 to 150 sq. ft. Repeat application in 4 to 6 weeks when necessary. Can be used as a supplement to Curme for control of Chicken Red Mites and Fleas. |
| POULTRY HOUSES | Northern Fowl Mites, Chicken Red Mites, Poultry Lice, Flies | 1 1/4 oz. per 10 gal. water | Apply liberally to floor, walls, and other parts of nest and adjacent area. For flies, mix with water and use as frequent applications as necessary. |

U.S.D.A. Reg. No. 359-526

Port. 5C-1-120

Atlanta, Ga
Bound Brook, N. J.
Chicago, Ill.
N. Kansas City, Mo

Manufactured by

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

Fresno, Calif.
Pasadena, Tex.
Portland, Ore.
St. Paul, Minn.

