

PRODUCT NO.

**2150**

NET CONTENTS  
GALLONS

# CHIPCO® TURF HERBICIDE MCPP

Especially useful for the control of surface-creeping broadleaf weeds such as Clovers, Chickweeds, Stitchwort, Ground Ivy, and Knotweed on sports and ornamental turf, including golf greens, fairways, and tees; as well as on home lawns and other non-crop turf areas.

**ACTIVE INGREDIENT:**

Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid\* 32.8%  
INERT INGREDIENTS 67.2%

\*Equivalent to 22.0% 2-(2-methyl-4-chlorophenoxy) propionic acid or 2.0 lbs. MCPP acid per gallon.

## CAUTION

### Keep Out of Reach of Children

Avoid contact with skin, eyes, or clothing. Avoid inhaling spray mist. Keep out of reach of children. Avoid spray drift to susceptible plants as this product may injure tomatoes, beans, peas, and ornamentals, (coarse sprays are less likely to drift). Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use the same equipment for other purposes). Do not transport or store near seeds, fertilizers, insecticides, or fungicides. Do not use in or near greenhouses. Avoid contamination of water intended for irrigation of crops or domestic purposes. Protect from freezing. If allowed to freeze, thaw and remix before using. Do not use on St. Augustine Grass. Do not reuse container. Destroy when empty.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**GENERAL INFORMATION:** MCPP has proven to be a very useful herbicide for control of the surface-creeping or prostrate broadleaf weeds listed below. Although relatively slow in action, one of the advantages is the tolerance of fine turf grasses to Chipco Turf Herbicide MCPP. Can be used on most Bent varieties, but use caution on Seaside, Arlington, and Congressional Bents which may be injured.

### WEEDS CONTROLLED

- Common Chickweed (Stellaria media)
- Mouseear Chickweed (Cerastium vulgatum)
- Red Clover (Trifolium pratense)
- White Clover (Trifolium repens)
- Ground Ivy (Glechoma hederacea)
- Stitchwort (Stellaria graminea)
- Knotweed (Polygonum aviculare)

### DIRECTIONS FOR USE

APPLICATION RATE ON:	PER 1,000 SQ. FT.	PER ACRE
Established greens and fine turf lawns	1½-2 fl. ozs.	4-5 pints
Other established turf and fairways	2½-3 fl. ozs.	6-7 pints

Better weed control, especially of Knotweed has been obtained with weekly applications at half recommended rate. This is preferred, especially on Bentgrasses and aprons where there has been no previous experience.

**HOW TO APPLY:** Apply as a water-mixed spray in 20 to 40 gals. water per acre (4 gals. per 1,000 sq. ft.). Apply with boom type sprayer to give thorough coverage. Smaller areas may be treated with a knapsack sprayer. Do not mow for several days before or after application. Withhold sprinkling for 24 hours after application.

**WHEN TO APPLY:** For Knotweed apply in spring or early summer when weeds are young for best results. For other weeds listed, apply in spring or fall when weeds are growing actively.

**Caution:** Do not apply when turf is suffering from drought conditions or during very hot or very wet weather or damage to turf may result.

Local conditions may affect the use of this chemical. Consult State Agricultural Experimental Station or Extension Service weed and turf specialists for specific recommendations as to dosage and time of application and for control of other weeds.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Do NOT Store with Foodstuffs

EPA Reg. No. 359-339-AA  
EPA Est. 359-MO-1  
EPA Est. 359-OR-1  
PL-0274

Manufactured by  
**CHIPMAN DIVISION**  
OF  
**RHODIA INC.**  
SOMERSET, NEW JERSEY

