

PRODUCT NO.

2170

NET CONTENTS
GALLONS

CHIPCO® TURF KLEEN

For control of many broadleaf weeds on golf course turf,
lawns and similar recreational and ornamental turf areas.

ACTIVE INGREDIENTS:

Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid*	16.2%
Diethanolamine salt of 2,4-dichlorophenoxy-acetic acid*	16.1%

INERT INGREDIENTS	67.7%
--------------------------	-------

Equivalent to 10.93% of 2-(2-methyl-4-chlorophenoxy) propionic acid or 1.0 lbs. MCPP acid per gallon. Equivalent to 10.93% 2,4-dichlorophenoxyacetic acid or 1.0 lbs. 2,4-D per gallon.

CAUTION

Keep Out of Reach of Children

Avoid contact with skin, eyes, or clothing. Avoid inhaling spray mist. Avoid spray drift to susceptible plants as this product may injure grapes, tomatoes, beans, peas, other vegetables, legumes, and ornamentals (coarse sprays are less likely to drift). Because of danger of contamination, do not use same spray equipment for applying insecticides and fungicides. Do not reuse shipping containers; destroy by crushing and burying where possible. Do not transport with or store near seeds, fertilizers, insecticides, or fungicides. Do not use in or around greenhouses. Avoid contamination of water intended for irrigation or domestic purposes. Store at temperatures above 32°F. If allowed to freeze, thaw and remix before using.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

GENERAL INFORMATION: MCPP has proven useful for control of surface-creeping or prostrate broadleaf weeds and 2,4-D controls many other broadleaf weeds as well as Wild Onions. Together these two chemicals will control the common broadleaf weeds growing in turf. May be used on all types of turf where 2,4-D is recommended including Merion, Kentucky Bluegrass, Bermudagrass, Fescues, and Ryegrass. Once the weeds have been removed the bare areas should be reseeded and the entire turf area given a good fertilizer treatment to encourage the grasses. Bare areas may be reseeded 3 to 4 weeks after application. Note: MCPP acts more slowly than 2,4-D, so that some weeds will be controlled before others.

Do Not Store with Foodstuffs

CONTROL THESE SUSCEPTIBLE WEEDS

Buckhorn; other plantains	Yarrow	White Clover
Curly Dock	English Daisy*	Ground Ivy
Dandelion*	Common Chickweed	Stitchwort
Red Sorrel*	Mouseear Chickweed	Knotweed
Wild Garlic*	Red Clover	Speedwell
Wild Onion*		

DIRECTIONS FOR USE

AMOUNT TO APPLY	PER 1,000 SQ. FT.	PER ACRE
Low rate	3¾ fl. oz.	5 quarts
High Rate	4½ fl. oz.	6 quarts

*Use high rate for weeds marked with an asterisk in the above list.

MIXING AND APPLYING: Mix with water, using 2 to 4 gallons of water per 1,000 sq. ft. On an acreage basis apply in 20 to 40 gallons of water per acre. Apply by means of a sprayer or watering can to give uniform coverage. Repeat application may be necessary especially with heavy infestations of the less susceptible weeds indicated thus *.

WHEN TO APPLY: Apply in spring or fall when weeds are actively growing. Knotweed is controlled more easily in spring or early summer when weeds are young.

CAUTION: Do not apply when turf is suffering from drought conditions or during very hot weather or damage to grass may result. Do not mow for several days before or after application. Withhold sprinkling for 24 hours after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas. Do not use if you want clover in your turf. In some areas Dichondra, Bentgrass, Carpetgrass, Buffalograss, and St. Augustinegrass may be injured; therefore do not use on turf where these grasses predominate.

Local conditions may affect the use of this chemical. Consult State Agricultural Experiment Station or Extension Service weed and turf control specialists for specific recommendations as to dosage and time of application and for control of other weeds.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

LOT No.

EPA Reg. No. 359-548-AA
EPA Est. 359-MO-1

PL-1274

Manufactured by

RHODIA INC.
AGRICULTURAL DIVISION

SOMERSET, NEW JERSEY



2 11/11/71