

BEST AVAILABLE COPY

NET CONTENTS
GALLONS

B U C T R I L^R

PRODUCT NO.
156

For the control of certain broadleaf weeds in small grains (wheat and barley)

ACTIVE INGREDIENT:

Octanoic acid ester of 3,5-dibromo-4-hydroxybenzonitrile*..... 33.9%
INERT INGREDIENTS..... 66.1%

*Bromoxynil octanoate
equivalent to 23.31% of 3,5-dibromo-4-hydroxybenzonitrile (bromoxynil)
2.0 lbs. of bromoxynil per gallon

C A U T I O N

KEEP OUT OF REACH OF CHILDREN

Do not take internally. Avoid contact with skin, eyes or clothing. In case of contact, flush eyes with plenty of water; get medical attention. Wash skin with soap and water. Avoid breathing spray mist. Avoid spray drift onto neighboring crops and pastures. Do not store near seeds, fertilizers, insecticides or fungicides. Do not re-use shipping containers; destroy by crushing and burying where possible. Avoid contamination of water intended for irrigation or domestic purposes. Contamination of shallow fish-bearing waters at these recommended rates will kill fish.

Control These Weeds in the Seedling Stage

Blue (Purple) Mustard (*Chorispora tenella*)
Fiddleneck (*Amsinckia intermedia*)
Corn Gromwell (*Lithospermum arvense*)
Cow Cockle (*Saponaria vaccaria*)
Field Pennywort (*Thlaspi arvense*)
Green Smartweed (*Polygonum scabrum*)
Lambsquarters (*Chenopodium album*)
London Rocket (*Sisymbrium irio*)
Shepherdspurse (*Capsella bursa-pastoris*)
Silverleaf Nightshade (*Solanum elaeagnifolium*)
Tartary Buckwheat (*Fagopyrum tataricum*)
Tarweed (*Hemizonia* spp.)
Tumble (Jim Hill) Mustard (*Sisymbrium altissimum*)
Wild Buckwheat (*Polygonum convolvulus*)
Wild Mustard (*Brassica kaber*)

D I R E C T I O N S

When To Apply - Apply after weeds have germinated and while they are small for best results. On fall seeded grain crops, apply in fall or spring, depending on weed species present. On spring grain, apply when grain is from 2-leaf to jointing stage.

Local conditions may affect the use of this chemical. Consult State or local agricultural authorities or weed specialists for specific recommendations as to dosage and time of application and for control of other weeds.

Seller makes no warranty of any kind, express or implied, concerning the results of use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

Amount to Use - Use 1 1/2 pints per acre in the young seedling stage. If weeds are beyond the very young seedling stage, use 2 pints.

How to Apply - Mix required quantity of Buctril with water. For ground spraying equipment, apply 10 to 15 gallons of spray per acre. For aerial application, apply 5 to 10 gallons of spray per acre.

apply after weeds have germinated and while still small and when grain is from 2 leaf up to boot stage

Do NOT Store with Foodstuffs

Manufactured by

CHIPMAN CHEMICAL COMPANY, INC.

Burlingame, California

Bound Brook, N.J.
Chicago, Ill.

Pasadena, Tex.
Palo Alto, Calif.

N. Kansas City, Mo.
Portland, Ore.

Bessemer, Ala.
St. Paul, Minn.

USDA Reg. No. 359-X

BB 9/10/65

