

PRODUCT NO.

156

BUCTRIL

NET CONTENTS
GALLONS

For the control of certain broadleaf weeds in wheat and barley.

ACTIVE INGREDIENT

Octanoic acid ester of 3,5-dibromo-4-hydroxy-
benzonitrile* 33.9%

INERT INGREDIENTS 66.1%

*Bromoxynil octanoate equivalent to 23.31% of 3,5-dibromo-4-hydroxybenzonitrile (bromoxynil). Contains 2.0 lbs. of bromoxynil per gallon.

CAUTION: Keep Out of Reach of Children

Do not take internally. Avoid contact with skin, eyes or clothing. In case of contact, flush eyes with plenty of water, get medical attention. Wash skin with soap and water.

Avoid breathing spray mist.

Avoid spray drift onto neighboring crops and pastures. Do not store near seeds, fertilizers, insecticides or fungicides. Do not re-use shipping containers, destroy by crushing and burying where possible. Avoid contamination of water intended for irrigation or domestic purposes. Contamination of shallow fish-bearing waters at these recommended rates will kill fish.

Local conditions may affect the use of this chemical. If lower dosages are to be used, consult State or local agricultural authorities or weed specialists for specific recommendations as to dosage and time of application and for control of other weeds.

Seller makes no warranty of any kind, express or implied, concerning the results of use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.




USDA Reg. No. 359-564

Do NOT Store with Foodstuffs

Part

CONTROL THESE WEEDS IN THE SEEDLING STAGE

Blue (Purple) Mustard (*Chorispora tenella*)
Fiddleneck (*Amsinckia intermedia*)
Corn Gromwell (*Lithospermum arvense*)
Cow Cockle (*Saponaria vaccaria*)
Field Pennycress (*Thlaspi arvense*)
Green Smartweed (*Polygonum scabrum*)
Groundsel (*Senecio vulgaris*)
Lambsquarters (*Chenopodium album*)
London Rocket (*Sisymbrium irio*)
Shepherdspurse (*Capsella bursa-pastoris*)
Silverleaf Nightshade (*Solanum elaeagnifolium*)
Tartary Buckwheat (*Fagopyrum tataricum*)
Tarweed (*Hemizonia* spp.)
Tumble (Jim Hill) Mustard (*Sisymbrium altissimum*)
Wild Buckwheat (*Polygonum convolvulus*)
Wild Mustard (*Brassica kaber*)



DIRECTIONS

WHEN TO APPLY: Apply after weeds have germinated and while still small and when grain is from 2-leaf up to boot stage.

AMOUNT TO USE: Use 1 1/2 pints per acre for weeds in the young seedling stage. If weeds are beyond the very young seedling stage use 2 pints.

HOW TO APPLY: Mix required quantity of BUCTRIL with water. For ground spraying equipment, apply 10 to 15 gallons of spray per acre. For aerial applications, apply 5 to 10 gallons of spray per acre.

Manufactured by

CHIPMAN CHEMICAL COMPANY, INC.

BURLINGAME, CALIFORNIA

Bessemer, Ala. Bound Brook, N. J. Chicago, Ill. N. Kansas City, Mo. Palo Alto, Calif. Pasadena, Tex. Portland, Ore. St. Paul, Minn.

Received Ref. 500
PAB-ALP
OCT 1 1965

ACCEPTED

NOV 12 1965

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO 359-564