

PRODUCT NO.

156

BUCTRIL[®]

For the control of certain broadleaf weeds in wheat and barley.

ACTIVE INGREDIENT

Octadecyl ester of 3,5-dibromo-4-hydroxybenzonitrile

33.9%

INERT INGREDIENTS

66.1%

15 emulsion containing 23.31% of 3,5-dibromo-4-hydroxybenzonitrile. Contains 2.1% of bronopol as preservative.

CAUTION: Keep Out of Reach of Children

Do not take internally. Avoid contact with skin, eyes or clothing. In case of contact, flush eyes with plenty of water, get medical attention. Wash skin with soap and water.

Avoid breathing spray mist.

Avoid spray drift onto neighboring crops and pastures. Do not use on crops, except for fertilizers, insecticides or fungicides. Do not re-use empty containers; do not by or into or blowing where or

Contamination of water with this chemical may kill fish. Rinse spray equipment and containers and dispose of liquid wastes in a pit or trench located away from water supply.

Environmental conditions may affect the use of this chemical. Consult state agricultural extension or Experiment Station weed specialists for specific recommendations for weed problems.

For more information of any of the above, contact your local agricultural extension or Experiment Station weed specialists for specific recommendations for weed problems.

Do not mix with Fertilizer Emulsions

WHEN TO APPLY: Make application before the weeds are in the seedling stage. It is important that weeds are removed soon after they have emerged when they are still seedlings and before they become established by the crop. Do not apply when plants are at seed from lack of moisture.

Manufactured by

ACCEPTED

8-26-66

359-564

CHIPMAN CHEMICAL COMPANY, INC.

BURLINGAME, CALIFORNIA

Boston, Mass. Bound Brook, N.J. Chicago, Ill. Denver, Colo. Kansas City, Mo. New York, N.Y. Pasadena, Tex. Portland, Ore.

CONTROL THESE WEEDS IN THE SEEDLING STAGE

- Blue (Purple) Mustard (Chorispora tenella)
- Fiddleneck (Tarweed) (Amsinckia intermedia)
- Corn Gromwell (Lithospermum arvense)
- *Cow Cockle (Sagittaria vaccaria)
- Field Pennycress (Thlaspi arvense)
- Green Smartweed (Polygonum scabrum)
- *Groundsel (Senecio vulgaris)
- Lambsquarters (Chenopodium album)
- *London Rocket (Sisymbrium irio)
- Shepherdspurse (Capsella bursa-pastoris)
- Silverleaf Nightshade (Solanum elaeagnifolium)
- Tartary Buckwheat (Fagopyrum tataricum)
- Tarweed (Hemizonia spp.)
- *Tumble (Jim Hill) Mustard (Sisymbrium aegyptium)
- Wild Buckwheat (Polygonum convolvulus)
- *Wild Mustard (Brassica kaber)

*Less susceptible weed

DIRECTIONS

IMPORTANT: Spray weeds soon after emergence and to prevent competition to the crop.

WHEN TO APPLY: Apply after the weeds have still in seedling stage and before they are in the crop. Do not apply when plants are lack of moisture.

AMOUNT TO USE: Use 1-1 1/2 pints per acre for susceptible weeds. For the less susceptible weeds use 2 pints per acre.

HOW TO APPLY: Apply as a water mix spray contact weed killer and good coverage of essential.

FOR GROUND APPLICATION EQUIPMENT: Apply 15 gallon total spray per acre. Use a nozzle if necessary to get good coverage. Nozzle should be at least 2 ft. above the crop.

FOR AERIAL APPLICATION: Apply in a minimum of 15 gallon total spray per acre. The plane should fly above the crop to insure good coverage. Nozzle with flow must not exceed length of spray.

Approved by
State of California
August 26, 1966



GALLONS

AGE

a)

31a)

lium)

tissimum)

ice for good weed
crop.

emerged and while
become shaded by
stressed from

for the more sus-
e weeds marked (*)

y. BUCTRIL is a
the weeds is

in a minimum of
higher gallonage
le pressure should

mum of 10 gallons of
fly at a height
at weeds. Width of
ray boom.

St. Paul, Minn.

ACCEPTED

AUG 26 1966

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 359-564