

BUCTRIL

NET CONTENTS
5 GALLONS

For the control of certain broadleaf weeds in wheat and barley.

DIET

and ester of 2,4-dichlorophenoxyacetic acid

benzothiazole

NTS

ate equivalent of 2,4-dichlorophenoxyacetic acid

UTION: Keep Out of Reach of Children

erally. Avoid contact with skin, eyes, nose, mouth, and clothing. Wash thoroughly with plenty of water after use. Attention: Wash hands after use.

g spray mist

to water, to be used on crops and pastures. Do not store near insecticides or other poisons. Do not use for shipping containers, crushing and baling where possible. Avoid contact with food for animals and domestic purposes.

to water with the above information. Do not use on crops and pastures. Do not use for shipping containers, crushing and baling where possible. Avoid contact with food for animals and domestic purposes.

L THESE WEEDS IN THE SEEDLING STAGE

stand (Chenopodium album)

weed (Amaranthus retrofractus)

(Lithospermum erythrorhizon)

long (Amaranthus)

(Lithospermum)

ed (Rumex crispus)

(Cirsium vulgare)

Chenopodium album

Sisymbrium irio

(Capsella bursa-pastoris)

hede (Chenopodium album)

al (Eragrostis tatarica)

onia spp.)

Mustard (Sisymbrium irio)

Polygonum conv. (L.)

assica (L.)

le (Poa annua) (L.)

le (Poa annua) (L.)

ranther (L.)

nettle (L.)

or Pepperweed (L.)

tible weed

DIRECTIONS

IMPORTANT: Apply weed killer after emergence for good weed control and to prevent competition to the crop.

WHEN TO APPLY: When applied before the crop has reached the boot stage. It is important that weeds are sprayed soon after they have emerged, when they are still susceptible to the action of the spray. Do not spray the crop when it is in flower or when the crop is about to be harvested.

AMOUNT TO USE: Use 1 pint per acre for the control of the weeds. For the control of the weeds, use 2 pints per acre.

HOW TO APPLY: Apply with water in a spray. BUCTRIL is a contact weed killer and good coverage of the weeds is essential.

FOR GROUND APPLICATION EQUIPMENT: Apply at a minimum of 25 gallons total spray per acre. Use a higher coverage if necessary to get good coverage. Nozzle pressure should be at least 20 p.s.i.

FOR AERIAL APPLICATION: Apply at a minimum of 10 gallons of total spray per acre. The spray should be applied in a way to insure good coverage of weeds. Wind speed should not exceed length of spray boom.

Do not use on crops or pastures. Consult State agricultural extension or Experiment Station weed specialists for specific recommendations for local weed problems.

Seller makes no warranty of any kind, express or implied, concerning the results of use of this product. Buyer assumes all risk of use or handling whether or not labeled with "Caution".

Do NOT Store with Foodstuffs

Manufactured by

CHIPMAN CHEMICAL COMPANY, INC.

BURLINGAME, CALIFORNIA

Go Bound Brook, N.J.

Chicago, Ill.

Denver, Colo.

N. Kansas City, Mo.

Palo Alto, Calif.

Pasadena, Tex.

Portland, Ore.

St. Paul, Minn.

159 564

