

COPY "A"

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.


COPY "B"

Newly Planted Grasses for Sod or Seed Production

Apply $1\frac{1}{2}$ pints of BUCTRIL per acre after grasses emerge and before broadleaf weeds are past the 3 to 4-leaf stage.

Use 2 pints of BUCTRIL per acre if weeds are beyond the 3 to 4 leaf stage but less than 6 inches in height and before flower formation. Weeds that form rosettes should be treated when these rosettes are less than $1\frac{1}{2}$ inches across.

This treatment may be used on seedling grasses such as Merion, Park, Delta or common Kentucky bluegrasses, Pennlawn, Chewings, Ulaker or Alta fescues, orchard grass, Highland, Seaside or Astoria bentgrasses, perennial ryegrasses planted for seed or sod production, or forage grasses planted for seed production purposes only.




PRODUCT NO.

156

BUCTRIL[®]

For the control of certain broadleaf weeds in wheat and barley



NET CONT 5 GAL

ACTIVE INGREDIENT

Octanoic acid ester of bromoxynil* (3,5-dibromo-4-hydroxybenzotrile)

33.9%

INERT INGREDIENTS

66.1%

*Bromoxynil octanoate equivalent is 23.31% of bromoxynil or not less than 20 pounds of bromoxynil per gallon.

CAUTION: Keep Out of Reach of Children

Avoid contact with skin, eyes or clothing. In case of contact, flush eyes with plenty of water, get medical attention. Wash skin with soap and water.

Avoid breathing spray mist

Avoid spray drift onto neighboring crops and pastures. Do not store near seeds, fertilizers, insecticides or fungicides. Do not re-use shipping containers; destroy by crushing and burying where possible. Avoid contamination of water intended for irrigation or domestic purposes.

Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands located away from water supply.

DIRECTIONS

IMPORTANT: Spray weeds soon after emergence for good weed control and to prevent competition to the crop.

WHEN TO APPLY: Make application before the crop has reached the boot stage. It is important that weeds are sprayed soon after they have emerged when they are still seedlings and before they become covered by the crop. Do not apply when plants are stressed from lack of moisture.

AMOUNT TO USE: Use 1 1/2 pints per acre for the more susceptible weeds. For the less susceptible weeds marked with an asterisk use 2 pints per acre.

HOW TO APPLY: Apply as a water mix spray. BUCTRIL is a contact weed killer and good coverage of the weeds is essential.

FOR GROUND APPLICATION EQUIPMENT: Apply in a minimum of 25 gallons total spray per acre. Use a higher gallonage if necessary to get good coverage. Nozzle pressure should be at least 20 p.s.i.

FOR AERIAL APPLICATION: Apply in a minimum of 10 gallons of total spray per acre. The plane should fly at a height above the crop to insure good coverage of weeds. Width of swath flown must not exceed length of spray boom.

CONTROL THESE WEEDS IN THE SEEDLING STAGE

Blue (Purple) Mustard (*Chorispora tenella*)

Corn Chamomile (Mayweed) (*Anthemis arvensis*)

Corn Gromwell (*Lithospermum arvense*)

*Cow Cockle (*Saponaria vaccaria*)

Fiddleneck (Tarweed) (*Amsinckia*)

Field Pennycress (*Thlaspi arvense*)

Green Smartweed (*Polygonum scrubrum*)

*Groundsel (*Senecio vulgaris*)

*Henbit (Dead Nettle) (*Lamium amplexicaule*)

Knawel (*Scrophanthus annuus*)

Lambsquarters (*Chenopodium album*)

London Rocket (*Sisymbrium irio*)

Peppergrass or Pepperweed (*Lepidium spp.*)

Shepherdspurse (*Capsella bursa-pastoris*)

Silverleaf Nightshade (*Solanum*)

Tarnary Buckwheat (*Fagopyrum tataricum*)

Tarweed (*Hemizonia spp.*)

*Tumble (Jim Hill) Mustard (*Sisymbrium altissimum*)

Wild Buckwheat (*Polygonum convolvulus*)

*Wild Mustard (*Brassica kaber*)

*Less susceptible weed.

→ COPY "5" HERE

Local conditions may affect the use of this chemical. Consult State or local extension or Experiment Station weed specialists for specific recommendations for local weed problems.

Follow directions carefully. Timing and method of application, weather, crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of material not in strict accordance with directions given herewith.

Do NOT Store with Foodstuffs

U.S.D.A. Reg. No. 359-564

Part. [redacted]

Atlanta, Ga.
Bound Brook, N. J.
Chicago, Ill.
St. Kansas City, Mo.

Manufactured by
RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

Palo Alto, Calif.
Pasadena, Tex.
Portland, Ore.
St. Paul, Minn.



HAZARDOUS IF SWALLOWED.

HAZARDOUS IF SWALLOWED.

SPERMICIDE

Mayweed (Dogfennel) (*Anthemis cotula*)

lycopsoide

elaegnifo