

156

ACCEPTED

4-16-68

5. 2 191 41 1 44

359-564

BUCTRIL

For the control of certain broadleaf weeds in wheat and barley

NET CONTENTS
5 GALLONS

ACTIVE INGREDIENT

Octanoic acid ester of promoxyr * 3,5-dibromo-4'-hydroxybenzonitrile 33.9%.

NEXT INGREDIENTS

5512.

*Breast cancer's equivalent to 23.3% of bromoxipr's not less than 5 pounds of bromoxipr per gallon

CAUTION: Keep Out of Reach of Children

Wash face with soap and water. Avoid contact with skin, eyes or clothing. In case of contact flush eyes with plenty of water, get medical attention. Wash skin with soap and water.

Avoid breathing spray mist

Avoid spraying drift onto neighboring crops and pastures. Do not store near seeds, fertilizers, insecticides or fungicides. Do not reuse in shipping containers, destroy by burning and burying where possible. Avoid contamination of water intended for irrigation or domestic purposes.

For a study equipment, the contents of the above of that cases in a p...
the goods ordered over the 1981-1982...

The product is for use as a...
The product is for use as a...
The product is for use as a...

CONTROL THESE WEEDS IN THE SEEDLING STAGE

6. Purple Mustard
 Brassica sp. (Brassicaceae)
 7. Orange Mustard
 Brassica sp. (Brassicaceae)
 8. Yellow Mustard
 Brassica sp. (Brassicaceae)
 9. Green Mustard
 Brassica sp. (Brassicaceae)
 10. Black Mustard
 Brassica sp. (Brassicaceae)
 11. White Mustard
 Brassica sp. (Brassicaceae)
 12. Red Mustard
 Brassica sp. (Brassicaceae)
 13. Blue Mustard
 Brassica sp. (Brassicaceae)
 14. Purple Mustard
 Brassica sp. (Brassicaceae)
 15. Orange Mustard
 Brassica sp. (Brassicaceae)
 16. Yellow Mustard
 Brassica sp. (Brassicaceae)
 17. Green Mustard
 Brassica sp. (Brassicaceae)
 18. Black Mustard
 Brassica sp. (Brassicaceae)
 19. White Mustard
 Brassica sp. (Brassicaceae)
 20. Red Mustard
 Brassica sp. (Brassicaceae)
 21. Blue Mustard
 Brassica sp. (Brassicaceae)

^aLess susceptible weed.

- *Lion's Rock's *Symbium* sp.
- Marweed *Daglene*
- Antennaria* sp.
- Peppergrass or Pepperweed
- Lesqueris* spp.
- Shepherdspurse
- Capsella bursa-pastoris*
- Silvered Nightshade
- Solanum elaeagnifolium*
- Tartary Buckwheat
- Fagopyrum tartaricum*
- Tarweed *Diemazium* spp.
- *Tumbleweed *M. Mustard*
- Symbium* sp.
- Wild Buckwheat
- Elymus canadensis*
- *W. M. yarrow *Erigeron* spp.

DIRECTIONS

IMPORTANT Spray weeds soon after emergence for good weed control and to prevent competition to the crop.

WHEN TO APPLY Make application before the crop has reached the boot stage. It is important that weeds are sprayed soon after they have emerged when they are still seedlings and before they become shaded by the crop. Do not apply when plants are stressed from lack of moisture.

AMOUNT TO USE Use 1/2 pint per acre for the more susceptible weeds. For the less susceptible weeds = seed in the 1st use 2 pints per acre.

HOW TO APPLY: Apply to water in a spray. BUCIPIL is a contact weed killer and good coverage of the weeds is essential.

FOR GROUND APPLICATION EQUIPMENT: Apply a minimum of 15 gallons total spray per acre. Use higher gallonage if necessary to get good coverage. Nozzle pressure 15-25 p.s.i.

FOR AERIAL APPLICATION: Apply at a rate of 5 gallons of total spray per acre. The spray should be applied to the top of the crop to insure good coverage and to reach the underside of the leaves. For best results, apply in the early morning or late afternoon.

NEWLY PLANTED GRASSES FOR SOD OR SEED PRODUCTION

Age: 1.5 years 13.0 before offer grossed emerge and before product

use 2 parts of B, CTF, then 4 parts of A, stage 2 is achieved with

The treatment was based on seedling grasses such as *Meibomia*, *Eragrostis*, and *Digitaria*, which are common in the Fens, by plowing the Fens down, growing a crop of *Antennaria* for two years, and then plowing the Fens down again. After the grasses, there are a few grasses, and a few years of seedling grasses, and a few grasses planted for seed production.

Users are encouraged to use the use of this material for personal study or for extension or experimentation. No need spending for special recommendations, for or weed control.

Following are the objectives of the study:

5. Various methods have been suggested for the storage of the material that is to be examined. The following are the most common:

Do NOT Store with Foodstuffs

54 Day
319 364

Atlanta Georgia
 Board of Directors
 Atlanta Georgia
 Atlanta Georgia

49-375-1316

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

[illegible]