


PRODUCT NAME: Buctril

REGISTRANT: RHODIA, INC., CHIPMAN DIVISION
LABEL AND REGISTRATION SECTION
120 JERSEY AVENUE, P.O. BOX 2009
NEW BRUNSWICK, NEW JERSEY 08903



WITH APPLICATION DATED: May 27, 1970

135

BUCTRIL[®]

For the control of certain broadleaf weeds in wheat () barley, oats, rye

ACTIVE INGREDIENT:

Octanoic acid ester of bromoxynil* (3,5-dibromo-4-hydroxybenzonitrile) 33.9%

INERT INGREDIENTS 66.1%

*Bromoxynil octanoate equivalent to 23.31% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact, flush eyes with plenty of water; get medical attention. Wash skin with soap and water.

Avoid breathing spray mist.

Avoid spray drift onto neighboring crops and pastures. Do not store near seeds, fertilizers, insecticides or fungicides. Do not re-use shipping containers; destroy by crushing and burying where possible. Avoid contamination of water intended for irrigation or domestic purposes.

Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands located away from water supply.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

CONTROL THESE WEEDS IN THE SEEDLING STAGE

Blue (Purple) Mustard
(*Chlorispora tenella*)
Corn Chenopod (Mayweed)
(*Anthemis arvensis*)
Corn Groundsail (*Lithospermum arvense*)
*Cow Chick (Saponaria vaccaria)
Field Chick (Tarweed)
(*Amsinckia intermedia*)
Field Pennywort (*Thlaspi arvense*)
Green Shepherd's Purse
(*Clethra racemosa*)
*Groundsail (*Senecio vulgaris*)
*Henbit (Dead Nettle)
(*Lamium amplexicaule*)
Knawel (*Scleranthus annus*)
Lambquarters (*Chenopodium album*)

*Less susceptible weed

*London Rocket (*Sisymbrium irio*)
Mayweed (Dogfennel)
(*Anthemis cotula*)
Peppergrass or Pepperweed
(*Lepidium spp.*)
Shepherdspurse
(*Capsella bursa-pastoris*)
Silverchick (Liquorice)
(*Solanum elaeagnifolium*)
Tatary Buckwheat
(*Fagopyrum tataricum*)
Tarweed (*Hemizonia spp.*)
*Tumble (Jim Hill) Mustard
(*Sisymbrium altissimum*)
Wild Buckwheat
(*Polygonum convolvulus*)
*Wild Mustard (*Brassica kaber*)

DIRECTIONS

IMPORTANT: Spray weeds soon after emergence for best results. Do not spray to prevent competition to the crop.

WHEN TO APPLY: Make application before the crop is well established. It is important that weeds are sprayed soon after they are still seedlings and before they become shaded by the crop. Plants are stressed from lack of moisture.

AMOUNT TO USE: Use 1 1/2 pints per acre for the most susceptible weeds listed in the list, use 2 pints for less susceptible weeds.

HOW TO APPLY: Apply as a water mix spray. BUCTRIL requires good coverage of the weeds is essential.

FOR GROUND APPLICATION EQUIPMENT: Apply 1 1/2 pints per acre. Use a higher gallonage if necessary. Nozzle pressure should be at least 20 p.s.i.

FOR AERIAL APPLICATION: Apply in a minimum of 100 gallons per acre. The plane should fly at a height above the crop of 100 feet. Width of swath flown must not exceed length of swath.

NEWLY PLANTED GRASSES FOR SOD OR

Apply 1 1/2 pints of BUCTRIL per acre after grasses emerge. Weeds are past the 3 to 4 leaf stage.

Use 2 pints of BUCTRIL per acre if weeds are beyond the 4 to 6 inch stage and before flower formation. Weeds should be treated when these rosettes are less than 1 inch high.

This treatment may be used on seedling grasses such as non-Kentucky bluegrasses, Pennlawn, Chewings, Illinoise, grass, Highland, Seaside or Astoria bentgrasses, perennials, seed or sod production, or forage grasses planted for sod only.

Local conditions may affect the use of this chemical. Consult Extension or Experiment Station weed specialists for suggestions on local weed problems.

Follow directions carefully. Timing and method of application, mixtures with other chemicals not specified, and other influencing factors in the use of this product are beyond the control of Bayer. Bayer assumes all risks of use, storage or handling in accordance with directions given here-with.

Do NOT Store with Foods

U.S.D.A. Reg.
No. 359-564

Atlanta, Ga.
Bound Brook, N. J.
Chicago, Ill.

Manufactured by

RHODIA INC.
CHIPMAN DIVISION

Palo Alto, Calif.
Pasadena, Tex.
Portland, Ore.
St. Louis, Mo.

