

PRODUCT NUMBER
2070

SUPPLEMENTAL LABELING
(EPA Reg. No. 359-564)

1-2087
BUCTRIL® HERBICIDE
RHÔNE-POULENC

WEED CONTROL IN COVER CROPS ON CONSERVATION RESERVE ACRES

DIRECTIONS FOR USE

CROPS:

Crested wheatgrasses (all), tall wheatgrass, western wheatgrass, bluebunch wheatgrass, intermediate wheatgrass, perennial ryegrass, fescues, bentgrasses, Kentucky bluegrass, Canby bluegrass, bahiagrass, Russian wildrye, seedling alfalfa

TARGET WEEDS:

BUCTRIL will control broadleaf weeds mentioned in the General Weed list found on the package label. Follow all use directions on this label and on the BUCTRIL package label to ensure product performance and crop safety.

WHEN TO APPLY:

BUCTRIL may be applied with standard ground boom and aerial hydraulic spray systems to cover crops listed here anytime after the 2-3 leaf stage, but before weeds exceed recommended stages of growth for treatment. Seeding alfalfa may be treated anytime after the 3 trifoliate leaf stage.

HOW TO APPLY

GROUND APPLICATION: Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with flat fan nozzles spaced a maximum of 20 inches apart and screens no finer than 50 mesh in the nozzle tips and in line strainers. Apply in a minimum spray volume of 10 gallons per acre with minimum spray pressure of 30 psi at the nozzle tips.

AERIAL APPLICATION: The sprayer should be equipped to provide a uniform and accurate application. Use a minimum spray volume of 5 gallons per acre using diaphragm type nozzles that produce cone or fan spray patterns with a maximum spray pressure of 40 psi at the nozzle tips. Nozzles should be located no further than 1/2 the distance from the center of the aircraft to the end of the wing or rotor. Nozzles should be oriented at an angle between straight down and straight back. The aircraft should discharge the spray a maximum of 10 feet above the crop. To minimize drift, do not apply during periods of gusty winds or when wind exceeds 5 MPH.

In order to ensure maximum safety and weed control, follow later recommendations on this label and all the cautions and limitations on the package label and of the tank mixture components.



RHÔNE-POULENC INC.
AGROCHEMICAL DIVISION

USE RECOMMENDATIONS

TIMING OF APPLICATION

RATE	CROP	WEED ^{1/}
1 1/2 Pints/A	Apply anytime in the fall or spring after grasses have reached the 2-3 leaf stage or when alfalfa has reached the 3-trifoliate leaf stage	MOST SUSCEPTIBLE BROADLEAF WEEDS Apply to weeds up to the 8 leaf stage or 4 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
2 Pints/A		SUSCEPTIBLE BROADLEAF WEEDS Apply to weeds up to the 4 leaf stage or 2 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.

^{1/} Refer to General Weed List on the Buctril label for a complete list of susceptible weeds

BUCTRIL AND MCPA

Grass Crops Only - Do not use on alfalfa

A tank mixture of BUCTRIL at the recommended rate + MCPA at 1/4 to 1/2 pound of active ingredient per acre is used to broaden the spectrum of broadleaf weed control. The tank mixture will control the weeds listed on the MCPA label in addition to the weeds listed on the BUCTRIL label. The rate of BUCTRIL in the tank mixture is the same as when BUCTRIL is used alone. See BUCTRIL RECOMMENDATIONS.

BUCTRIL + MCPA tank mixtures can be applied after the 3 leaf stage but before the crop reaches the boot stage. Apply only to the following grass species: Crested wheatgrasses, tall wheatgrass, bluebunch wheatgrass, intermediate wheatgrass, sheep fescue, Canby bluegrass, Russian wildrye. Do not apply to alfalfa.

USE RESTRICTIONS:

Do not allow livestock to graze treated fields.

Do not feed treated grasses or alfalfa to livestock.

Do not apply BUCTRIL + MCPA tank mixtures to alfalfa.

Mix with clean water only. do not use spray additives or fluid fertilizers as crop injury may occur.

Do not apply BUCTRIL to alfalfa when air temperatures exceed 80° F as crop injury may occur.

This labeling must be in the possession of the user at the time of post-harvest application.

All appropriate directions, restrictions, and precautions on the BUCTRIL and MCPA package labels are to be followed.

In order to ensure maximum safety and weed control, follow label recommendations on this label and all the cautions and limitations on the package labels of the tank mixture components.

USE RECOMMENDATIONS:

BUCTRIL

TIMING OF APPLICATION

RATE	CROP	WEED ^{1/}
1 1/2 Pints/A	Apply anytime in the fall or spring after grasses have reached the 2-3 leaf stage or when alfalfa has reached the 3-trifoliate leaf stage.	Most Susceptible Broadleaf Weeds. Apply to weeds up to the 8 leaf stage or 4 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
2 Pints/A		Susceptible Broadleaf Weeds. Apply to weeds up to the 4 leaf stage or 2 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.

¹ Refer to the general weed list on the Bucril package label for a complete list of susceptible weeds.

BUCTRIL AND MCPA - Grass Crops Only - Do not use on alfalfa.

A tank mixture of Bucril at the recommended rate + MCPA at 1/4 to 1/2 pound of active ingredient per acre is used to broaden the spectrum of broadleaf weed control. The tank mixture will control the weeds listed on the MCPA label. In addition, to the weeds listed on the Bucril label. The rate of Bucril in the tank mixture is the same as when Bucril is used alone. See BUCTRIL RECOMMENDATIONS.

Bucril + MCPA tank mixtures can be applied after the 3 leaf stage but before the crop reaches the boot stage. Apply only to the following grass species Crested wheatgrasses, tall wheatgrass, blue bunch wheatgrass, intermediate wheatgrass, sheep fescue, canby bluegrass, Russian wildrye. Do not apply to alfalfa.

USE RESTRICTIONS:

Do not allow livestock to graze treated fields.

Do not feed treated grasses or alfalfa to livestock.

Do not apply Bucril + MCPA tank-mixtures to alfalfa.

Mix with clean water only, do not use spray additives or fluid fertilizers as crop injury may occur.

Do not apply Bucril to alfalfa when air temperatures exceed 80°F as crop injury may occur.

This labeling must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, and precautions on the Bucril and MCPA package labels are to be followed.

