



## BUCTRIL HERBICIDE SEEDLING ALFALFA USE DIRECTIONS

### SEEDLING ALFALFA

Apply BUCTRIL when weed seedlings are actively growing. For optimum control apply BUCTRIL in sufficient spray volume to ensure good coverage of the weeds.

### BUCTRIL RECOMMENDATIONS

BUCTRIL can be applied to seedling alfalfa with a minimum of 2 trifoliate leaves.

Apply BUCTRIL at 1 pint/A to control the MOST SUSCEPTIBLE BROADLEAF WEEDS and 1 1/2 pints/A for control of SUSCEPTIBLE BROADLEAF WEEDS (see GENERAL WEED LIST). Weeds should not exceed the 4 leaf stage or 2 inches in height, whichever comes first, or before rosettes are 1.5 inches in diameter. For improved control of kochia, pigweed (sp), tansymustard and henbit BUCTRIL at 1 pint/A should be applied in tank mixture with 2,4-DB at 0.5 ai/A. BUCTRIL will not adequately control over-wintered pennycress, henbit and mustards.

For ground application use a minimum of 20 gallons of spray volume per acre. Use flat fan nozzles spaced a maximum of 20 inches apart with a minimum spray pressure of 30 psi at the nozzle tips. For aerial application use a minimum of 5 gallons of spray volume per acre with a maximum spray pressure of 40 psi at the nozzle tips. Refer to AERIAL APPLICATION under GENERAL INFORMATION section of the BUCTRIL label for complete use directions. For use through automated sprinkler irrigation systems BUCTRIL should be applied at 2 pints per acre only. Refer to the GENERAL INFORMATION Section of the BUCTRIL label for complete sprinkler irrigation system use directions.

### RESTRICTIONS AND PRECAUTIONS

Crop leaf burn can occur following BUCTRIL application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected. Alfalfa yield should not be reduced although total biomass tonnage may decrease compared to a weedy field due to weed removal.

Do not treat when temperatures exceed 70° F at and 3 days following application because unacceptable crop injury may occur.



RESTRICTIONS AND PRECAUTIONS (cont.)

Do not apply when alfalfa is under moisture, temperature, insect or disease stress.

Do not add a surfactant or crop oil because increased alfalfa injury will occur.

Do not cut for feed or graze spring treated alfalfa within 30 days following treatment.

Do not cut for feed or graze fall or winter treated alfalfa until spring, at least 60 days following treatment

Total BUCTRIL applied to a seedling alfalfa crop should not exceed 2 pints/A.

Follow all restrictions and precautions on the 2,4-DB label when a BUCTRIL + 2,4-DB tank mixture is used.

Do not apply BUCTRIL through automated sprinkler irrigation systems in California.

Do not apply BUCTRIL + 2,4-DB tank mixture through automated sprinkler irrigation systems.

