

PRODUCT NUMBER
2038

SUPPLEMENTAL LABELING
(EPA Reg. No. 359-601)

Label No. 2038
1-20-87

BRONATE[®] HERBICIDE



WEED CONTROL IN COVER CROPS ON CONSERVATION RESERVE ACRES

DIRECTIONS FOR USE

CROPS:

Crested wheatgrass, smooth wheatgrass, weath wheatgrass, bluebunch wheatgrass, intermediate wheatgrass, sideoats grama, and sandbar. Also, weedy and giant ragweed.

TARGET PESTS:

BRONATE will control a broad range of weeds listed on the General Weed List found on the package label. Follow all use directions on the label and on the BRONATE package label to ensure product performance and crop safety.

WHEN TO APPLY:


BRONATE may be applied with standard ground boom and air hydraulic spray systems to cover crops listed here anytime after the 2-leaf stage, but before weeds reach recommended stages of growth for treatment. Refer to Use Recommendations below for further entry dates.

HOW TO APPLY:

GROUND APPLICATION: Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with flat fan nozzles spaced a maximum of 20 inches apart and screens or filter finer than 50 mesh in the nozzle tips and in line strainers. Apply in a minimum spray volume of 10 gallons per acre with minimum spray pressure of 30 psi at the nozzle tips.

AERIAL APPLICATION: The sprayer should be equipped to provide a uniform and accurate application. Use a minimum spray volume of 5 gallons per acre ground equivalent. Use nozzles that produce cone or fan spray patterns with a maximum spray pressure of 40 psi at the nozzle tips. Nozzles should be located no further than 1/2 the distance from the center of the aircraft to the end of the wing or rotor. Nozzles should be oriented at an angle between straight down and straight back. The aircraft should discharge the spray a maximum of 10 feet above the crop. Do not apply during periods of gusty winds or when wind exceeds 5 MPH.

In order to ensure maximum safety and weed control, follow label recommendations on this label and all the cautions and limitations on the package labels of the tank mixture components.



RHÔNE-POULENC INC.
AGROCHEMICAL DIVISION

USE RECOMMENDATIONS

TIMING OF APPLICATION

BRONATE

RATE	CROP	WEED ^{1/}
1 1/2 Pints A	Apply anytime in the fall or spring after the crop has reached the 2-3 leaf stage	MOST SUSCEPTIBLE BROADLEAF WEEDS. Apply to weeds up to the 8 leaf stage or 4 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
2 Pints A		SUSCEPTIBLE BROADLEAF WEEDS. Apply to weeds up to the 4 leaf stage or 2 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter. Apply to henbit, knawel, and mayweed up to the 4 leaf stage or 2 inches in height, whichever comes first.

^{1/} Refer to General Weed List on the Bronate label for a complete list of susceptible weeds.

USE RESTRICTIONS:

Do not allow livestock to graze treated fields.

Do not feed treated grasses to livestock.

Mix with clean water only; do not use spray adjuvants or surfactants as crop injury may occur.

Do not apply to Alfalfa.

This labeling must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

PRODUCT NUMBER
2038

SUPPLEMENTAL LABELING

BRONATE^R HERBICIDE

WEED CONTROL IN COVER CROPS
ON CONSERVATION RESERVE ACRES
DIRECTIONS FOR USE

CROPS:

Crested wheatgrasses (all), tall wheatgrass, western wheatgrass, bluebunch wheatgrass, intermediate wheatgrass, sheep fescue Russian wildrye and Canby bluegrass.

TARGET PESTS:

Bronate will control all broadleaf weeds mentioned in the General Weed list found on the package label. Follow all use directions on this label and on the Bronate package label to ensure product performance, and crop safety.

WHEN TO APPLY:

Bronate may be applied with standard ground boom and aerial hydraulic spray systems to forage crops listed here anytime after the 2-3 leaf stage, but before weeds exceed recommended stages of growth for treatment. Refer to Use Recommendations below for further instructions.

HOW TO APPLY:

Ground Application: Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with flat fan nozzles spaced a maximum of 20 inches apart and screens no finer than 50 mesh in the nozzle tips and in-line strainers. Apply in a minimum spray volume of 10 gallons per acre with minimum spray pressure of 30 psi at the nozzle tips.

Aerial Application: The sprayer should be equipped to provide a uniform and accurate application. Use a minimum spray volume of 5 gallons per acre using diaphragm-type nozzles that produce cone or fan spray patterns with a maximum spray pressure of 40 psi at the nozzle tips. Nozzles should be located no further than 3/4 the distance from the center of the aircraft to the end of the wing or rotor. Nozzles should be oriented at an angle between straight down and straight back. The aircraft should discharge the spray a maximum of 10 feet above the crop. To minimize drift, do not apply during periods of gusty winds of when wind exceeds 5 MPH.

In order to ensure maximum safety and weed control, follow label recommendations on this label and all the cautions and limitations on the package labels of the tank mixture components.

USE RECOMMENDATIONS

BRONATE

TIMING OF APPLICATION

RATE	CROP	WEED ^{1/}
1 1/2 Pints/A	Apply anytime in the fall or spring after the crop has reached the 2-3 leaf stage.	Most Susceptible Broadleaf Weeds. Apply to weeds up to the 8 leaf stage or 4 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
2 Pints/A		<p>Susceptible Broad eaf Weeds. Apply to weeds up to the 4 leaf stage or 2 inches in height whichever come first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.</p> <p>Apply to henbit, knawal, and mayweed up to the 4 leaf stage or 2 inches in height, whichever comes first.</p>

^{1/} Refer to General Weed List on the Bronate label for a complete list of susceptible weeds.

USE RESTRICTIONS:

Do not allow livestock to graze treated fields.

Do not feed treated grasses to livestock.

Mix with clean water only, do not use spray adjuvants or fluid fertilizers as crop injury may occur.

Do not apply to Alfalfa.

