

VISKO-RHAP formulations are designed to produce thick invert emulsions when used in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water enter the spray system separately to be mixed in a special chamber as the spraying is performed.

WHERE TO USE

VISKO-RHAP LOW VOLATILE 2TP controls certain woody and herbaceous plants in rights-of-way, industrial sites, fencerows, and similar noncrop areas.

PLANTS CONTROLLED

VISKO-RHAP LOW VOLATILE 2TP will kill or control the following woody and herbaceous plants as well as others less susceptible to 2,4-D or 2,4,5-T.

WOODY GROWTH

Blackjack oak	Poison ivy	Scrub oak
Ground-ivy	Post oak	White oak
Mesquite	Red oak	Wild blackberry
Pin oak	Salt cedar	Yucca
	Sand shinnery oak	

EMERGENT AQUATIC WEEDS

Alligatorweed	Pickeralweed	Spatterdock
Arrowhead	Rush	Waterlily

DIRECTIONS

This special formulation is designed for use in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

If this product is to be applied through a bifluid nozzle, sodium bisulfate water-conditioning agent must be added at the rate of 2 pounds per 50 gallons of water.

Best results are obtained when spraying is done while plants are in a rapid stage of growth and after leaves are fully developed. Repeat applications may be necessary if new growth appears.

Application rates of 1 to 2 gallons of product per acre are recommended depending on plants density, height, and species.

Aerial Application: Fill water tank with fresh, clean water. Add VISKO-RHAP LOW VOLATILE 2TP herbicide to the formulation tank. Use settings to give desired chemical-to-water ratio in the range of 1:3 to 1:7. Adjust pressure to suit conditions, but do not exceed 40 p.s.i.

For Emergent Aquatic Weeds in lakes, ponds, drainage ditches, marshes, and similar water areas: Use 1 to 2 gallons of product per acre to control Arrowhead, Pickeralweed, Rush, Spatterdock and Waterlily; use 2 to 4 gallons of product per acre to control Alligatorweed. Application should be made when leaves are fully developed above the water line and plants are actively growing. DO NOT apply to more than 1/2 to 1/2 of a lake or pond in any one month, because excessive decaying vegetation may deplete oxygen content of water, killing fish.

PRODUCT NO.

709-A



NET CONTENTS
GALLONS

WARNING

Avoid spray drift to susceptible broad-leaved plants that are desirable such as cotton, soy beans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

This product when used as directed forms a thick invert emulsion that breaks up into large spray droplets, which are least likely to drift. However, minute quantities can cause injury by drift, and application conditions that permit drift of spray should be avoided. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

DO NOT graze dairy animals on treated areas within seven days after application.

DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose. DO NOT apply insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of the product.

DO NOT reuse empty containers. Destroy by burying in an isolated place away from water supplies.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. In case of contact with skin or eyes, immediately flush with plenty of water and get medical attention for eyes. Wash clothing before reuse. Keep away from heat or open flame.

Keep out of reach of children.

Follow directions carefully. Timing and method of application, weather conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions give herewith.

DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME

VISKO-RHAP

LOW VOLATILE ESTER

2TP

HERBICIDE

FOR AIR APPLICATION

2,4,5-TP

2 LBS OF SILVEX EQUIVALENT PER GALLON

ACCEPTED

3-23-72

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR RECORD NO. 359-644

FOR AIR APPLICATION

CAUTION: KEEP OUT OF REACH OF CHILDREN.

READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

The product in this container is especially formulated and recommended for use in the process of U. S. Patent 3,197,299. In consideration of the payment of royalty of \$0.75 per gallon of product in this container Rhodia Inc., Chipman Division hereby grants to the purchaser of said product a non-exclusive license to practice the process of U. S. Patent 3,197,299 with said product. Licenses to practice said process with material from any source are available at the same royalty from Rhodia Inc., Chipman Division, New Brunswick, N. J. 08903.

ACTIVE INGREDIENT:	
2-Ethylhexyl Ester of Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	34.4%
(Equivalent to 24.33% Silvex) (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	
INERT INGREDIENTS	65.6%
†An Isooctyl Ester	TOTAL 100.0%

Manufactured by

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

EPA Reg. No. 359-644-AA
LICENSED FOR USE UNDER U. S. PATENT NO. 3,197,430

PRINTED IN U.S.A.

Port. 15M-1-1271