

VISKO-RHAP formulations are designed to produce thick invert emulsions when used in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water enter the spray system separately to be mixed in a special chamber as the spraying is performed.

WHERE TO USE

VISKO-RHAP LOW VOLATILE 2D is used to control broad-leaved weeds in rights-of-way, industrial sites, fence-rows, similar noncrop areas; and for aquatic weed control in drainage ditches, ponds, lakes, marshes, and similar waters.

WEEDS CONTROLLED

VISKO-RHAP LOW VOLATILE 2D will kill or control the following weeds as well as many others susceptible to 2,4-D:

| | | |
|-----------------------------|----------------|-----------------|
| Alligatorweed | Curly indigo | Purslane |
| Arrowhead | Duckweed | Rush |
| Artichoke | Elderoerry | Russian |
| Bindweed | Goldenrod | thistle |
| (Hedge, field and European) | Ground ivy | Sagebrush |
| Bitter | Hemp | Shepherds-purse |
| wintergrass | Hoary cress | Smartweed |
| Boxelder | Honeysuckle | Sowthistle |
| Buckhorn | Indigo | Spatterdock |
| Bull thistle | Ironweed | Stinkweed |
| Burdock | Jimsonweed | Sumac |
| Bur ragweed | Lambs-quarters | Sunflower |
| Buttercup | Locoweed | Virginia |
| Canada | Mexicanweed | creeper |
| thistle | Morningglory | Waterhyacinth |
| Catnip | Mustard | Waterlily |
| Chickweed | Pickeralweed | Water |
| Chicory | Pigweed | smartweed |
| Cocklebur | Plantain | Wild garlic |
| Coffeebean | Poison ivy | Wild lettuce |
| Creeping | Pokeweed | Wild onion |
| Jenny | Povertyweed | Wild radish |
| | Puncturevine | Willow |

DIRECTIONS

This special formulation is designed for use in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

If this product is to be applied through a bifluid nozzle, sodium bisulfate water-conditioning agent must be added at the rate of 2 pounds per 50 gallons of water.

Best results are obtained when spraying is done while weeds are young and actively growing.

Application rates of 2 to 6 pounds per acre (1 to 3 gallons of product of 2,4-D acid equivalent are recommended depending on weed density, height, and species).

For emergent and floating aquatic weeds in drainage ditches, lakes, ponds, marshes and similar water areas: Use 1 to 2 gallons of product per acre to control the emergent weeds Arrowhead, Pickerelweed, Rush, Spatterdock, Waterlily and Water smartweed; use 2 to 3 gallons of product per acre to control the emergent weed Alligatorweed. Use 1 gallon of product per acre to control the floating weed Duckweed. Application should be made when leaves are fully developed above water line and plants are actively growing. Depending on plant size use sufficient volume of spray to give coverage of all weeds. Repeat application if regrowth occurs. Consult your State Game and Fish Department or Water Control Agency prior to application of this product for aquatic weed control.

DO NOT apply to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen content of water, killing fish.


Ground Application (Handgun, Manifold, or Spray Boom): Fill water tank with fresh, clean water. Add VISKO-RHAP Low Volatile 2D herbicide to the formulation tank. Use orifices to give desired chemical to water ratio in the range of 17 to 115. Adjust pressure to suit the spray job being done. Do not exceed 40 p.s.i. when using spray boom. Do not exceed 100 p.s.i. when using handgun or manifold.

PRODUCT NO.

708-G



NET CONTENTS
GALLONS



WARNING

Avoid spray drift to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

This product when used as directed forms a thick invert emulsion that breaks up into large spray droplets, which are least likely to drift. However, minute quantities can cause injury by drift, and application conditions that permit drift of spray should be avoided. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

DO NOT apply this product on lawns. Legumes, such as white clover, are susceptible and may be severely damaged or killed.

DO NOT graze dairy animals on treated areas within seven days after application.

DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose. DO NOT spray insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of the product.

DO NOT reuse empty containers. Destroy by burying in an isolated place away from water supplies.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. In case of contact with skin or eyes, immediately flush with plenty of water and get medical attention for eyes. Wash clothing before reuse. Keep away from heat or open flame.

Keep out of reach of children.

Follow directions carefully. Timing and method of application, weather conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

VISKO-RHAP

LOW VOLATILE ESTER

2D

HERBICIDE

FOR GROUND APPLICATION

2,4-D

2 LBS. OF 2,4-D ACID EQUIVALENT PER GALLON

CAUTION: KEEP OUT OF REACH OF CHILDREN.

READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

The product in this container is especially formulated and recommended for use in the process of U. S. Patent 3,197,299. In consideration of the payment of royalty of \$0.75 per gallon of product in this container Rhodia Inc., Chipman Division hereby grants to the purchaser of said product a non-exclusive license to practice the process of U. S. Patent 3,197,299 with said product. Licenses to practice said process with material from any source are available at the same royalty from Rhodia Inc., Chipman Division, New Brunswick, N. J. 08903.

| | |
|--|---------------|
| ACTIVE INGREDIENT: | |
| 2-Ethylhexyl Ester† of 2,4-Dichlorophenoxyacetic Acid‡ | 37.1% |
| INERT INGREDIENTS: | |
| | 62.9% |
| TOTAL | 100.0% |
| †An Isooctyl Ester | |
| ‡Equivalent to 24.6% of 2,4-Dichlorophenoxyacetic Acid | |

Manufactured by

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY