

VISKO-RHAP formulations are designed to produce thick invert emulsions when used in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water enter the spray system separately to be mixed in a special chamber as the spraying is performed.

WHERE TO USE

VISKO-RHAP A-3T controls certain woody and herbaceous plants in rights-of-way, industrial sites, fencerows, and similar noncrop areas.

PLANTS CONTROLLED

VISKO-RHAP A-3T will kill or control the following as well as other noxious plants susceptible to 2,4,5-T:

Ash	Maple
Aspen	Mesquite
Buckbrush	Osageorange
Chokecherry	Poison ivy
Cuttant	Poison oak
Elder	Poplar
Hawthorn	Scrub oak
Hickory	Wild blackberry

DIRECTIONS

This special formulation is designed for use in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

If this product is to be applied through a bifluid nozzle, sodium bisulfate water-conditioning agent must be added at the rate of 2 pounds per 50 gallons of water.

Best results are obtained when spraying is done while weeds are in a rapid stage of growth and after leaves are fully developed. Soil moisture should be adequate for good growth and it is preferable that relative humidity be high. Repeat applications may be necessary if new growth appears.

Application rates of 1 quart to 2 gallons of VISKO-RHAP A-3T per acre are suggested depending on weed or brush density, height, and species. Prior to placing VISKO-RHAP A-3T in the supply tank, it should be diluted and mixed thoroughly with at least 1 part chemical to 1 part No. 2 fuel oil or kerosene and can be diluted with up to 3 parts oil to each part of chemical. This will enable the operator to control rate and volume of application.

Aerial Application: Fill water tank with fresh, clean water. Dilute and add VISKO-RHAP A-3T to the chemical tank. Use settings to give desired chemical-to-water ratio in the range of 1:3 to 1:7. Adjust pressure to suit the job being done. Do not exceed 40 p.s.i. at the nozzles.

Ground Application: Dilute product and fill tanks as directed above for aerial application. Use orifices to give desired chemical to water ratio in the range of 1:5 to 1:15. Adjust pressure to suit conditions. Do not exceed 40 p.s.i. with boom. Do not exceed 100 p.s.i. with handgun or manifold.

PRODUCT NO.
713



NET CONTENTS
GALLONS
ACCEPTED

VISKO-RHAP

**OIL-SOLUBLE AMINE
A-3T
HERBICIDE**

2,4,5-T

3 LBS OF 2,4,5-T ACID EQUIVALENT PER GALLON

CAUTION: KEEP OUT OF REACH OF CHILDREN.

READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

The product in this container is especially formulated and recommended for use in the process of U. S. Patent 3,197,299. In consideration of the payment of royalty of \$0.75 per gallon of product in this container Rhodia Inc., Chipman Division hereby grants to the purchaser of said product a non-exclusive license to practice the process of U. S. Patent 3,197,299 with said product. Licenses to practice said process with material from any source are available at the same royalty from Rhodia Inc., Chipman Division, New Brunswick, N. J. 03903

ACTIVE INGREDIENT:	
N,N-Dimethyl Oleyl-Linoleyl Amine Salt of 2,4,5-Trichloro- phenoxyacetic Acid (Equivalent to 34.84% 2,4,5-T Acid)	57.0%
INERT INGREDIENTS:	43.0%
TOTAL	100.0%

Manufactured by
RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

WARNING

DO NOT apply this product to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

This product when used as directed forms a thick invert emulsion that breaks up into large spray droplets, which are least likely to drift. However, minute quantities can cause injury by drift, and application conditions that permit drift of spray should be avoided.

DO NOT graze dairy animals on treated areas within 6 weeks after application.

DO NOT graze meat animals on treated areas within 2 weeks of slaughter.

DO NOT contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. This product is toxic to fish.

DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. DO NOT contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose. DO NOT apply insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of the product.

DO NOT reuse empty containers. Destroy by burying in an isolated place away from water supplies.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. In case of contact with skin or eyes, immediately flush with plenty of water and get medical attention for eyes. Wash clothing before reuse. Keep away from heat or open flame.

Keep out of reach of children.

Follow directions carefully. Timing and method of application, weather conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME**

**DO NOT USE AROUND THE HOME
OR RECREATION AREAS**