

Visko-Rhap formulations are designed to produce thick invert emulsions when used in the Chipman Division of Rhodia Inc. Two-Fluid Spray System. The formulation and water enter the spray system separately to be mixed in a special chamber as the spraying is performed.

### WHERE TO USE

Visko-Rhap A-3D controls weeds in rice, small grains, and grain sorghum; and certain woody and herbaceous plants in rights-of-way, industrial sites, fencerows, and similar noncrop areas.

### PLANTS CONTROLLED

Visko-Rhap A-3D will kill or control the following as well as other noxious plants susceptible to 2,4-D:

Blackberry	Dryland Willow	Mexicanweed	Sand Shinnery
Bur Ropeweed	Duckweed	Oak	Sunflower
Cockspur	Hawthorn	Perennial	Tea Tree
Cottonweed	Hickory	Ragweed	Willow
Curly Indigo	Indigo	Sagebrush	Waterhyacinth
	McCartney Rose	Salt Cedar	
	Mesquite		

### DIRECTIONS

This special formulation is designed for use in the Chipman Division of Rhodia Inc., Two-Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

If this product is to be applied through a bi-fluid nozzle, sodium bisulfate water-conditioning agent must be added at the rate of 2 pounds per 50 gallons of water.

Annual and perennial weeds are most easily killed in the young seedling stage. Established perennials are best controlled at the time flower buds appear. Woody species are most susceptible after leaves are fully developed and when plants are actively growing.

**WATERHYACINTH CONTROL** in still water (lakes, ponds, and marshes): Application rate of 1 gallon of Visko-Rhap A-3D per acre is suggested. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted and thoroughly mixed with one part chemical to one part No. 2 fuel oil or kerosene. Boat Application: See Ground Application recommendations. Consult your State Game and Fish Department or Water Control Agency prior to application of this product for aquatic weed control.

### GENERAL WEED AND BRUSH CONTROL

Application rate of 1 quart to 1 gallon of Visko-Rhap A-3D per acre is suggested, depending on weed density, height, and species. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted and mixed thoroughly with at least 1 part chemical to 1 part No. 2 fuel oil or kerosene, and can be diluted with up to 3 parts oil to each part of chemical. This will enable the operator to control rate and volume of the spray.

### WEED CONTROL IN CROPS

**DILUTION:** When using Visko-Rhap A-3D as a weed control agent in crops, it should be diluted as directed above.

**RICE:** Treat 6 to 9 weeks after seeding with ½ to 1 gallon properly diluted Visko-Rhap A-3D per acre in control of Curly Indigo, Indigo, and Mexicanweed.

**SMALL GRAINS (wheat, barley, rye):** Spray when grain is in full tiller stage (4-5 inches tall) with ½ to 1 gallon properly diluted Visko-Rhap A-3D per acre.

Do not use in grain intended for legumes, unless damage to legume can be tolerated. Do not spray small grain after boot stage.

Do not forage or graze treated grain fields within 2 weeks after treatment.

Do not feed treated straw to livestock.

**GRAIN SORGHUMS:** Treat when grain is 6 to 12 inches tall and a secondary root system is well developed with ½ to 1 gallon properly diluted Visko-Rhap A-3D per acre. Use dropped nozzles when grain is over 10 inches high. Best results are obtained if spraying is done when weeds are in a rapid stage of growth.

**Aerial Application:** Fill water tank with fresh, clean water. Dilute and add Visko-Rhap A-3D to the chemical tank. Use settings to give desired chemical-to-water ratio in the range of 1:3 to 1:7. Adjust pressure to suit the job being done. Do not exceed 40 p.s.i. at the nozzles.

**Ground Application:** Dilute product and fill tanks as directed above for aerial application. Use orifices to give desired chemical-to-water ratio in the range of 1:5 to 1:15. Adjust pressure to suit conditions. Do not exceed 40 p.s.i. with boom. Do not exceed 100 p.s.i. with handgun or manifold.

PRODUCT NO.

2905



NET CONTENTS  
GALLONS

# VISKO-RHAP

## OIL-SOLUBLE AMINE

ACCEPTED

MAY 24 1974

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC PESTS REGISTERED  
UNDER NO. 359-650-1

## A-3D

## HERBICIDE

## 2,4-D

3 LBS OF 2,4 D ACID EQUIVALENT PER GALLON

### CAUTION: KEEP OUT OF REACH OF CHILDREN

READ ENTIRE LABEL FOR ADDITIONAL HANDLING PRECAUTIONS.

By virtue of an agreement with the patent owner, the purchaser of the material in this container is hereby authorized to practice with said material the processes of U. S. Patents 3,197,299; 3,592,385; and 3,640,461. Similar authorization to practice the processes of said patents with material from any source is available from the patent owner.

<b>ACTIVE INGREDIENT:</b>	
n,n-dimethyl oleyl-oleyl amine salt of 2,4-dichloro- phenoxyacetic acid	63.86%
(Equivalent to 35.46% 2,4-D Acid)	
<b>INERT INGREDIENTS:</b>	36.14%
<b>TOTAL</b>	<b>100.00%</b>

Manufactured by

**CHIPMAN DIVISION  
OF  
RHODIA INC.  
SOMERSET, NEW JERSEY**

EPA Reg. No. 359-650-AA  
EPA Est. 359-MO-1  
LICENSED FOR USE UNDER U. S. PATENT 2,930,411

### WARNING

Do not apply this product to susceptible broadleaf plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

This product when used as directed forms a thick invert emulsion that breaks up into large spray droplets, which are least likely to drift. However, minute quantities can cause injury by drift, and application conditions that permit drift of spray should be avoided.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. This product is toxic to fish. Do not contaminate water used for irrigation or domestic purposes.

Do not graze dairy animals on treated pastures within seven days after application.

Do not store this product near fertilizers, seeds, insecticides, or fungicides. Do not contaminate feed or foodstuffs.

Do not use spray equipment contaminated with this product for any other purpose. Do not apply insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

Do not mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40° F. and agitate before using. This does not affect the efficiency of the product.

Do not reuse empty containers. Destroy by burying in an isolated place away from water supplies.

### CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. In case of contact with skin or eyes, immediately flush with plenty of water and get medical attention for eyes. Wash clothing before reuse. Keep away from heat or open flame.

Keep out of reach of children.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given here-with.

**DO NOT USE, POUR, SPILL, OR STORE  
NEAR HEAT OR OPEN FLAME**

LOT NO.