


visko-rhap

HERBICIDE

Oil-Soluble
Amine A-30

18



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Keep this and all other pesticides out of the reach of children. Do not use this product near food, feed, or food containers. Do not use this product on food crops or animals.

ENVIRONMENTAL HAZARDS

Keep this and all other pesticides out of the reach of children. Do not use this product near food, feed, or food containers. Do not use this product on food crops or animals.

DIRECTIONS FOR USE

Read label carefully. Use only as directed. Do not use on food crops or animals.

STORAGE AND DISPOSAL

Store in original container. Do not use if container is damaged. Do not use if contents are solidified. Do not use if contents are discolored. Do not use if contents are contaminated. Do not use if contents are expired. Do not use if contents are otherwise unsuitable for use.

PESTICIDE DISPOSAL

Pesticide spray mixture should not be used on food crops or animals. Do not use if contents are solidified. Do not use if contents are discolored. Do not use if contents are contaminated. Do not use if contents are expired. Do not use if contents are otherwise unsuitable for use.

CONTAINER DISPOSAL

Do not reuse container for food, feed, or food containers. Do not use if container is damaged. Do not use if contents are solidified. Do not use if contents are discolored. Do not use if contents are contaminated. Do not use if contents are expired. Do not use if contents are otherwise unsuitable for use.

Always use pesticides as directed on the label.

For more information, contact your local extension office.

NOTE

Read label carefully. Use only as directed. Do not use on food crops or animals. Do not use if container is damaged. Do not use if contents are solidified. Do not use if contents are discolored. Do not use if contents are contaminated. Do not use if contents are expired. Do not use if contents are otherwise unsuitable for use.

WHERE TO USE

Visko-Rhap A-3D controls weeds in rice, sorghum, and indigo, sorghum, and cotton, woody and herbaceous plants, rights of way, and other areas where weeds and brush are a problem.

PLANTS CONTROLLED

Visko-Rhap A-3D will control the following as well as other susceptible plants:

Blackberry	Dryland Weeds	Meadow weed	Sand Spurry
Blue Ragweed	Duckweed	" "	Sumac
Cocklebur	Hawthorn	Persimmon	Sunflower
Coffeebean	Hickory	Rabbitbrush	Tan yarrow
Cottonwood	Indigo	Ragweed	Willow
Curly Indigo	Macartney Rose	Sagebrush	Waterhyacinth
	Mesquite	Salt Cedar	

This special formulation is designed for use in the Rhine-Potomac Chemical Company Two Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

Annual and perennial weeds are most easily killed in the young seedling stage. Established perennials are best controlled at the time flower buds appear. Woody species are most susceptible after leaves are fully developed and when plants are actively growing.

WATERHYACINTH CONTROL - Fresh water, lake, ponds and marshes.

Application rate of 1 gallon of Visko-Rhap A-3D per acre is suggested. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted 1 part to 10 parts with one part chemical to one part No. 2 fuel oil or kerosene. Boat Application: See Ground Application recommendations. Consult your State Game and Fish Department or Water Control Agency prior to application of this product for aquatic weed control.

GENERAL WEED AND BRUSH CONTROL

Application rate of 1 part to 1 gallon of Visko-Rhap A-3D per acre is suggested, depending on weed density, height, and species. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted and mixed thoroughly with at least 1 part chemical to 1 part No. 2 fuel oil or kerosene, and can be diluted with up to 3 parts of 1 part of chemical. This will enable the operator to control rate and volume of the spray.

WEED CONTROL IN CROPS

DILUTION

When using Visko-Rhap A-3D in water, it should be diluted as directed.

RICE

1 gallon of Visko-Rhap A-3D per acre is suggested. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted and mixed thoroughly with at least 1 part chemical to 1 part No. 2 fuel oil or kerosene, and can be diluted with up to 3 parts of 1 part of chemical. This will enable the operator to control rate and volume of the spray.

SMALL GRAINS

1 gallon of Visko-Rhap A-3D per acre is suggested. Prior to placing Visko-Rhap A-3D in the supply tank, it should be diluted and mixed thoroughly with at least 1 part chemical to 1 part No. 2 fuel oil or kerosene, and can be diluted with up to 3 parts of 1 part of chemical. This will enable the operator to control rate and volume of the spray.

Treat when grain is 6-12 inches tall and secondary root systems well developed with 1 to 1 1/2" galler properly diluted VIKSJO RHAP-A 3D per acre. Use dropped nozzles when grain is over 10 inches high. Best results are obtained if spraying is done when wheat is at the boot or early stage of growth.

Ground Application: Dilute product in 100 times as much water as needed for aerial application. Use nozzles to give desired rate to water ratio. The standard rate is 1.0 fl. oz. per 100 gal. Adjust pressure to suit application. Do not exceed 40 psi at nozzle. Do not use equipment with a discharge manifold.

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The *Agrobacterium* strains were grown in the YEA medium for 24 h at 28 °C. The cell concentration of the strains was adjusted to 1.0 × 10⁸ cells/ml. The cell suspension was mixed with the plant tissue and the transformation efficiency was determined. The results were expressed as the mean ± SD of three independent experiments. The asterisks indicate the significant difference between the strains at the same concentration of the cell suspension.