

Visko-Rhap formulations are designed to produce thick invert emulsions when used in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water enter the spray system separately to be mixed in a special chamber as the spraying is performed.

#### WHERE TO USE

Visko-Rhap Low Volatile 2DP is used to control broad-leaved weeds in rights-of-way, industrial sites, fencerows, drainage ditchbanks and fire-breaks; and similar non-crop areas.

#### WEEDS CONTROLLED

Visko-Rhap Low Volatile 2DP will control Sand sagebrush and Shinnery oak.

#### DIRECTIONS

This special formulation is designed for use in the Rhodia Inc., Chipman Division Two-Fluid Spray System. The formulation and water SHOULD NOT be mixed before use.

If this product is to be applied through a bifluid nozzle, sodium bisulfate water-conditioning agent must be added at the rate of 2 pounds per 50 gallons of water.

Best results are obtained when spraying is done while weeds are young and actively growing.

Application rates of 1/2 to 1 pound per acre of 2,4-DP acid equivalent (1 to 2 quarts of formulation) are recommended depending on brush density and height. Spray to cover foliage of unwanted vegetation. Two successive applications spaced a year apart are recommended for Shinnery oak.

**Aerial Application:** Fill water tank with fresh, clean water. Water temperature should not be less than 55°F. Dissolve sodium bisulfate water-conditioning agent (supplied with formulation) at the rate of 2 pounds per 50 gallons of water. Add Visko-Rhap Low Volatile 2DP herbicide to the formulation tank. Use orifices giving (formulation:water) ratio in the range of 1:3 to 1:9. Adjust pressure regulator on the pumps to suit the spray job being done, but do not exceed 35 p.s.i.


PRODUCT NO.

722



NET CONTENTS  
GALLONS

# VISKO-RHAP



**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

READ SIDE PANELS FOR ADDITIONAL HANDLING PRECAUTIONS.

The product in this container is especially formulated and recommended for use in the process of U. S. Patent 3,197,299. In consideration of the payment of royalty of \$0.75 per gallon of product in this container Rhodia Inc., Chipman Division hereby grants to the purchaser of said product a non-exclusive license to practice the process of U. S. Patent 3,197,299 with said product. Licenses to practice said process with material from any source are available at the same royalty from Rhodia Inc., Chipman Division, New Brunswick, N. J. 08903.

<b>ACTIVE INGREDIENT:</b>	
Isocetyl ester of 2-(2,4-dichlorophenoxy) propionic acid*	36.3%
*Equivalent to 24.6% 2,4-DP acid or not less than 2.0 lbs. of 2,4-DP acid per gallon.	
<b>INERT INGREDIENTS</b>	<b>63.7%</b>
<b>TOTAL</b>	<b>100.0%</b>

Manufactured by  
**RHODIA INC.**  
CHIPMAN DIVISION  
NEW BRUNSWICK, NEW JERSEY

EPA Reg. No. 359-654  
LICENSED FOR USE UNDER U. S. PATENT NO. 3,197,299

#### WARNING

Avoid spray drift to susceptible broad-leaved plants that are desirable such as cotton, soybeans, tomatoes, grapes, clover, ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

This product when used as directed forms a thick invert emulsion that breaks up into large spray droplets, which are least likely to drift. However, minute quantities can cause injury by drift, and application conditions that permit drift of spray should be avoided. Although this product contains an ester of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

**DO NOT** contaminate water in irrigation ditches or water used for domestic purposes.

**DO NOT** apply this product on lawns. Legumes, such as white clover, are susceptible and may be severely damaged or killed.

**DO NOT** graze dairy animals on treated areas within 6 weeks after application.

**DO NOT** graze meat animals on treated areas within 2 weeks of slaughter.

**DO NOT** store this product near fertilizers, seeds, insecticides, or fungicides. **DO NOT** contaminate feed or foodstuffs.

**DO NOT** use spray equipment contaminated with this product for any other purpose. **DO NOT** spray insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

**DO NOT** mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40°F. and agitate before using. This does not affect the efficiency of this product.

**DO NOT** reuse empty containers. Destroy by burying in an isolated place away from water supplies.

#### CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. In case of contact with skin or eyes, immediately flush with plenty of water and get medical attention for eyes. Wash clothing before reuse. Keep away from heat or open flame.

The water solution of sodium bisulfate used with this product to form the invert emulsion is slightly acid, and should not be used for any other purpose. It may be corrosive to metals and clothing or irritating to the skin. If splashed on the skin or in eyes, wash with plenty of water and get medical attention for eyes. Splashes on skin, clothes, or other materials may be neutralized with baking soda.

Keep out of reach of children.

Follow directions carefully. Timing and method of application, weather conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**DO NOT USE AROUND HOME OR RECREATION AREAS**

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

PRINTED IN U.S.A.

Part. 2C-1-0372