

Product Number 3355

# CHIPCO<sup>®</sup> 26019

## FUNGICIDE

A Foliar Applied Fungicide for the Prevention and Control of Certain Diseases of Turfgrass

### ACTIVE INGREDIENT:

3-(3,5-dichlorophenyl)-N-(1-methylethyl)  
-2,4-dioxo-1-imidazolidinecarboxamide ..... 50.0%

INERT INGREDIENTS ..... 50.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

See Back Panel for Additional Precautionary Statements



Manufactured By

**rhône-poulenc inc.**

agricultural division

monmouth junction, n.j. 08852

### NET CONTENTS

2 POUNDS  
(.907 Kg)

EPA Reg. No. 259-685  
EPA Est. No.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get Medical Attention if irritation persists.

### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

#### GENERAL USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage, disposal, cleaning of equipment. Do not store near other pesticides or seeds. Open dumping is prohibited.

**CONTAINER DISPOSAL:** Dispose of in a landfill approved for pesticide containers, or bury in a safe place.

Consult Federal, State or Local Disposal Authorities for approved alternative procedures.

### GENERAL INFORMATION

Chipco 26019 is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, home lawns, and institutional areas where fine turf is grown.

When used in conjunction with good turf management practices, Chipco 26019 is effective in the following diseases:

**Summer Diseases**—DOLLAR SPOT, BROWN PATCH and FUSARIUM BLIGHT on all cool season grasses. HELMINTHOSPORIUM LEAFSPOT and MELTING OUT is controlled on Bluegrass, and Bermudagrass.

157

157

2

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get Medical Attention if irritation persists.

**ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

**GENERAL USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds. Open dumping is prohibited.

**CONTAINER DISPOSAL:** Dispose of in a landfill approved for pesticide containers, or buy in a safe place.

Consult Federal, State or Local Disposal Authorities for approved alternative procedures.

**GENERAL INFORMATION**

Chipco 26019 is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, home lawns, and institutional areas where fine turf is grown.

When used in conjunction with good turf management practices, Chipco 26019 is effective in controlling the following diseases:

**Summer Diseases**—DOLLAR SPOT, BROWN PATCH and FUSARIUM BLIGHT on all common turf-grasses. HELMINTHOSPORIUM LEAFSPOT and MELTING OUT is controlled on Bluegrass, Bent-grass, and Bermudagrass.

157

157A

**Winter Diseases**—FUSARIUM PATCH (Pink Snow Mold) is controlled in the Pacific Northwest (the Cascade Mountains) only. Pink Snow Mold and Gray Snow Mold are suppressed in the Pacific Northwest.

Apply the recommended rates as indicated in the table below in 2 to 10 gallons of water. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

Disease	Interval of Applications	Oz
Dollar Spot ( <i>Sclerotinia homeocarpa</i> ) Brown Patch ( <i>Rhizoctonia solani</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	
Helminthosporium: Leaf Spot (Including Red Leaf Spot) Melting Out ( <i>Helminthosporium spp.</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	
Fusarium Blight ( <i>Fusarium roseum</i> )	Foliar applications should begin when conditions first become favorable for disease development. Two additional applications should be made at 2 to 3 week intervals.	
Fusarium Patch ( <i>Fusarium nivale</i> ) PACIFIC NORTHWEST ONLY (West of the Cascade Mountains)	Begin applications when conditions favor disease development. Repeat at 14-21 day intervals as long as required.	
SUPPRESSION ONLY: Gray Snow Mold ( <i>Typhula spp.</i> ) Pink Snow Mold ( <i>Fusarium nivale</i> )	One application before first permanent snow fall. If possible, another application during a midwinter thaw.	

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for these diseases. Do not mow or irrigate treated areas until the foliage is completely dry. A waiting period following treatment is preferred. Do not mix with any sticker, extender, or surfactant. Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock.

Local conditions may affect the use of this fungicide. Consult State Agricultural Experiment Station specialists for specific recommendations, consistent with label, for local use.

This product conforms to the chemical description on the label thereof and is registered on such label only when used in accordance with directions under normal conditions. Directions are given for use under normal conditions. Timing and method of application, weather and crop conditions are factors not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, handling, and disposal in strict accordance with directions given herewith.

**BEST DOCUMENT AVAILABLE**

2

**PRECAUTIONARY STATEMENTS  
DANGERS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

On skin, eyes, or clothing. In case of contact immediately wash with plenty of water. Get Medical Attention if irritation occurs.

**ENVIRONMENTAL HAZARDS**

Streams, and ponds. Do not contaminate water by cleaning or disposal of wastes.

**DIRECTIONS FOR USE**

**GENERAL USE**

Read entire label and directions of Federal Law to use this product in a manner consistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage, disposal or equipment. Do not store near other pesticides or seeds. Open flames prohibited.

**DISPOSAL:** Dispose of in a landfill approved for pesticide disposal or bury in a safe place.

Consult Federal, State or Local Disposal Authorities for approved alternatives.

**GENERAL INFORMATION**

Systemic fungicide, recommended for turfgrass disease control on golf courses, parks, and institutional areas where fine turf is grown.

Under good turf management practices, Chipco 26019 is effective in controlling

DOLLAR SPOT, BROWN PATCH and FUSARIUM BLIGHT on all common turfgrasses. FUSARIUM LEAFSPOT and MELTING OUT is controlled on bluegrass, Bentgrass.

**Winter Diseases**—FUSARIUM PAT (Pink snow mold) is controlled in the Pacific Northwest (West of the Cascade Mountains) only. Pink Snow Mold and Gray Snow Mold are suppressed in areas other than the Pacific Northwest.

Apply the recommended rates as indicated in the table below in 2 to 10 gallons of water per 1000 sq. ft. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

Disease	Interval of Applications	RATE
		Oz. Product/1000 ft <sup>2</sup>
Dollar Spot ( <i>Sclerotinia homeocarpa</i> ) Brown Patch ( <i>Rhizoctonia solani</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	1.5-2.0
Helminthosporium: Leaf Spot (Including Red Leaf Spot) Melting Out ( <i>Helminthosporium spp.</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	2.0
Fusarium Blight ( <i>Fusarium roseum</i> )	Foliar applications should begin when conditions first become favorable for disease development. Two additional applications should be made at 2 to 3 week intervals.	4.0
Fusarium Patch ( <i>Fusarium nivale</i> ) PACIFIC NORTHWEST ONLY (West of the Cascade Mountains)	Begin applications when conditions favor disease development. Repeat at 14-21 day intervals as long as required.	2.0-4.0
SUPPRESSION ONLY: Gray Snow Mold ( <i>Typhula spp.</i> ) Pink Snow Mold ( <i>Fusarium nivale</i> )	One application before first permanent snow fall. If possible, another application during a midwinter thaw.	2.0-4.0

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24 hour waiting period following treatment is preferred. Do not mix with any sticker, extender, or wetting agent. Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock or poultry.

Local conditions may affect the use of this fungicide. Consult State Agricultural Extension or Experiment Station specialists for specific recommendations, consistent with label, for local pest problems.

This product conforms to the chemical description on the label thereof and is reasonably fit for purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, handling of this material, not in strict accordance with directions given herewith.

**BEST DOCUMENT AVAILABLE**

PL-0180

157

157A

158

Pink #  
Mold

controlled in the Pacific Northwest (West of  
w Mold are suppressed in areas other than

in 2 to 10 gallons of water per 1000 sq. ft.  
as some breakdown of the product may  
a properly calibrated sprayer.

	RATE
	Oz. Product/1000 ft <sup>2</sup>
itions favor the disease first intervals as long	1.5-2.0
itions favor the disease first intervals as long	2.0
in when able for disease applications k intervals.	4.0
itions favor t at 14-21 day	2.0-4.0
ermanent snow cation during a	2.0-4.0

ival of applications are recommended for all  
oliage is completely dry, usually a 24 hour  
with any sticker, extender, or wetting agent.  
s from treated turf to livestock or poultry.

it State Agricultural Extension or Experiment  
it with label, for local pest problems.

bel thereof and is reasonably fit for purpose  
ecutions under normal use conditions. Follow  
ther and crop conditions, mixture with other  
cing factors in the use of this product are  
use, storage, handling of this material, not in

ILE

PL-0180

158

359-68

ACCEPTED  
20 FEB 1980  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
this pesticide was found to be  
effective under EPA Reg. No. 359-685.

1 0 0 0  
0 0 0 0