

PRODUCT NUMBER  
3355

**SPECIMEN LABEL**



**A Foliar Applied Fungicide for the Prevention and Control  
of Certain Diseases of Turfgrass**

ACTIVE INGREDIENT:  
3-(3,5-dichlorophenyl)-N-(1-methylethyl)  
-2,4-dioxo-1-imidazolidinecarboxamide ..... 50.0%  
INERT INGREDIENTS ..... 50.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
**STATEMENT OF PRACTICAL TREATMENT**  
In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes  
and get medical attention  
See Back Panel for Additional Precautionary Statements

Manufactured by  
**RHÔNE-POULENC CHEMICAL COMPANY**  
**AGROCHEMICAL DIVISION**  
**RHÔNE-POULENC INC.**  
Monmouth Junction, N J 08852

EPA Reg. No. 359-685  
EPA Est. No.  
NJ-0880

NET CONTENTS  
2 POUNDS  
.907 Kg

165

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get Medical Attention if irritation persists.

**ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

**GENERAL USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:**

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds. Open dumping is prohibited.

**CONTAINER DISPOSAL:**

Dispose of in a landfill approved for pesticide containers, or bury in a safe place.

Consult Federal, State or Local Disposal Authorities for approved alternative procedures.

359-685

166

## GENERAL INFORMATION

Chipco 26019 is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, home lawns, and institutional areas where fine turf is grown.

When used in conjunction with good turf management practices, Chipco 26019 is effective in controlling the following diseases:

**Summer Diseases —**

DOLLAR SPOT, BROWN PATCH and FUSARIUM BLIGHT on all common turfgrasses. HELMINTHOSPORIUM LEAFSPOT and MELTING OUT is controlled on Bluegrass, Bentgrass, and Bermudagrass.

**Winter Diseases —** FUSARIUM PATCH (Pink snow mold) is controlled in the Pacific Northwest (West of the Cascade Mountains) only. Pink Snow Mold and Gray Snow Mold are suppressed in areas other than the Pacific Northwest.

Apply the recommended rates as indicated in the table below in 2 to 10 gallons of water per 1000 sq. ft. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

Disease	Interval of Applications	RATE
		Oz. Product/ 1000 ft <sup>2</sup>
Dollar Spot ( <i>Sclerotinia homeocarpa</i> ) Brown Patch ( <i>Rhizoctonia solani</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	1.5-2.0
Helminthosporium: Leaf Spot (Including Red Leaf Spot) Melting Out ( <i>Helminthosporium spp.</i> )	Begin applications when conditions favor disease development or when the disease first appears. Repeat at 14-21 day intervals as long as required.	2.0
Fusarium Blight ( <i>Fusarium roseum</i> )	Foliar applications should begin when conditions first become favorable for disease development. Two additional applications should be made at 2 to 3 week intervals.	4.0
Fusarium Patch ( <i>Fusarium nivale</i> ) PACIFIC NORTHWEST ONLY (West of the Cascade Mountains)	Begin applications when conditions favor disease development. Repeat at 14-21 day intervals as long as required.	2.0-4.0
SUPPRESSION ONLY: Gray Snow Mold ( <i>Typhula spp.</i> ) Pink Snow Mold ( <i>Fusarium nivale</i> )	One application before first permanent snow fall. If possible, another application during a midwinter thaw.	2.0-4.0

Under severe conditions, the higher rate and/or shorter interval of applications are recommended. Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24 hour waiting period following treatment is preferred. Do not mix with any sticker, extender, or wetting agent. Do not graze animals on treated turf. Do not feed to animals on treated turf to livestock or poultry.

Local conditions may affect the use of this fungicide. Consult State Agricultural Extension or Experiment Station inspectors for specific recommendations, consistent with label, for local pest problems.

This product conforms to the chemical description on the label thereof and is reasonably fit for purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, handling of this material, not in strict accordance with directions given herewith.

359-685