

PRODUCT NUMBER

3357

Rowral



FUNGICIDE

ACCEPTED

AUG 27 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 359-685

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide	50.0%
INERT INGREDIENTS	50.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 359-685
 EPA Est. No.
 0682

NET CONTENTS
2 POUNDS
 .907 Kg

Manufactured by
RHÔNE-POULENC INC.
AGROCHEMICAL DIVISION
 Monmouth Junction, N.J. 08852

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get Medical Attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTION FOR USE

GENERAL USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

ACCEPTED

APR 27 1982

GENERAL INFORMATION

392-685

Rovral is a wettable powder formulation for the control of blossom and fruit monilinia brown rot on CHERRIES, PEACHES and NECTARINES. Apply as a foliar spray in sufficient water to obtain thorough coverage (20-400 gallons per acre by ground application and 15 to 20 gallons by aerial application.) The spray mixture should not be allowed to stand overnight as some breakdown of the product may occur from prolonged delays, particularly in water with a high pH. Maintain agitation during spray operations and apply with properly calibrated application equipment.

NOTE TO USER: Do not graze livestock in treated orchards. Do not feed cover crops grown in treated orchards to livestock.

Do not make more than 5 applications of Rovral per season.

App Rovral in accordance with directions provided in the table below.

DISEASE	Dosage Rate		SPRAY SCHEDULE
	Lbs. Prod/ 100 gal.	Lbs. Prod/ A	
Brown Rot Blossom Blight (<i>Monilinia</i> sp.)	0.25 to 0.50	1.0 to 2.0	Apply first at early bloom (Approximately 5% bloom). If conditions are favorable for disease development apply again at full bloom and at petal fall. 10/
Fruit Brown Rot (<i>Monilinia</i> sp.)	0.25 to 0.50	1.0 to 2.0	An application should be made whenever temperatures and moisture conditions favor disease infection in the five week period prior to harvest. If these conditions persist or reoccur, a second application should be made. This second application should be made no sooner than 7 days following the first preharvest application. Preharvest applications may be made up to and including the day of harvest.

Under severe disease conditions the higher rate is recommended.

The rates of Rovral per 100 gallons are based on a standard of 400 gallons per acre dilute spray for mature trees. For less than mature trees, apply the rate per 100 gallons until runoff. If less than 400 gallons of spray solution per acre is applied to mature trees, refer to the rate per acre to insure that the proper amount of material is applied.

Do not apply more than 2.0 lbs. of Rovral per acre per application.

CONDITION OF SALE, WARRANTY, LIMITATION OF LIABILITY

This product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of Rhone-Poulenc or the seller. Buyer assumes all risks of use, storage, or handling of this material, not in strict accordance with directions given herewith. In no case shall Rhone-Poulenc or the seller be liable for consequential, special, or indirect damages such as loss of profits or values resulting from the use or handling of this product.