


Product Number
3357

Proposed Label



ROVRAL®

Fungicide

Active Ingredient:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)
-2,4-dioxo-1-imidazolidinecarboximide.....50.02
Inert Ingredients.....50.02

Keep Out Of Reach Of Children

CAUTION

Statement of Practical Treatment

In case of contact, wash skin with soap and water; for eyes
flush with water for 15 minutes and get medical attention.

See Back Panel for Additional Precautionary Statements

BEST DOCUMENT AVAILABLE

EPA Reg. No. 359-685
EPA Est.

0983

Net Contents
1 Pounds
.907 Kg

Manufactured by

RHONE-POULENC INC
Agrochemical Division
Monmouth Junction, New Jersey 08852

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS**

CAUTION

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTION FOR USE

General Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEST DOCUMENT AVAILABLE

GENERAL INFORMATION

Rovral is a wettable powder formulation for the control of Brown Rot Blossom Blight and suppression of Shothole on almonds. Apply as a foliar spray in sufficient water to obtain thorough coverage (100-400 gallons per acre by ground application and 20 gallons by aerial application). The spray mixture should not be allowed to stand overnight as some breakdown of the product may occur from prolonged delays, particularly in water with a high pH. Maintain agitation during spray operations and apply with properly calibrated application equipment.

NOTE TO USER: Do not graze animals in treated orchards. Do not feed cover crops grown in treated orchards to livestock.

Do not make more than 2 applications of Rovral per season.

Apply Rovral in accordance with directions provided in the table below.

DISEASE	DOSAGE RATE		SPRAY SCHEDULE
	LBS. PRODUCT/ 100 GAL.	LBS. PROD./ ACRE	
Brown Rot Blossom Blight (<u>Monilinia</u> <u>laxa</u>)	0.25	1.0	Apply first at pink bud stage and again at full bloom when conditions are favorable for disease development.
SUPPRESSION: Shothole (<u>Coryneum</u> <u>beijerinckii</u>)			

The rates of Rovral per 100 gallons are based on a standard of 400 gallons per acre dilute spray for mature trees. For less than mature trees, apply the rate per 100 gallons until runoff. If less than 400 gallons of spray solution per acre is applied to mature trees, refer to the rate per acre to insure that the proper amount of material is applied.

CONDITION OF SALE, WARRANTY, LIMITATION OF LIABILITY

This product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of Rhone-Poulenc or the seller. Buyer assumes all risks of use, storage, or handling of this material, not in strict accordance with directions given herewith. In no case shall Rhone-Poulenc or the seller be liable for consequential, special, or indirect damages such as loss of profits or values resulting from the use or handling of this product.