


Product Number
5090

Ethoprop



(TECHNICAL GRADE)



For Formulation Into End Use Insecticide/Nematicide Products
Intended ONLY for Uses Accepted By the United States Environ-
mental Protection Agency.

ACTIVE INGREDIENT:

O-Ethyl-S, S-Dipropyl Phosphorodithioate94%
INERT INGREDIENTS6%

DANGER!  **POISON**
KEEP OUT OF REACH OF CHILDREN

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Give 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician, Poison Control Center, or hospital emergency room.

If not breathing, give mouth to mouth artificial respiration.

In Case of Skin Contact: Immediately remove all contaminated clothing, and wash skin with soap and water. Wash clothing before reuse. Discard contaminated shoes.

For Eyes: Flush with plenty of water for at least 15 minutes. Call a physician immediately in all cases of suspected poisoning.

Persons providing first aid should prevent contamination of themselves.

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 309-694
EPA EST. 359-TN-1
0183

NET CONTENTS
POUNDS
(Kilos)

Manufactured By

RHÔNE-POULENC INC.

AGROCHEMICAL DIVISION

Monmouth Junction, New Jersey 08852

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Wear waterproof protective clothing, rubber gloves and goggles. In case of contact, wash immediately with soap and water.

Do not breathe fumes or spray mist. Wear an AO R-6058 respirator with an R-58 cartridge, or equivalent, for protection during field handling and field exposures. After use, wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all clothing with soap and hot water before reuse. Discard contaminated shoes. Do not contaminate feed or food.

Do not store in or around the home.

SYMPTOMS OF POISONING: Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

NOTE TO PHYSICIAN: This product is a cholinesterase inhibitor. Severe symptoms and signs include diarrhea, pin-point and non-reactive pupils, respiratory difficulty, pulmonary edema, cyanosis, loss of sphincter control, convulsions, and coma.

Support respiration as needed. Measures should include removal of secretions, maintenance of a patent airway, and if necessary, artificial ventilation.

If cyanosis is absent, give ATROPINE 2-4 mg intravenously (0.05 mg/kg for children). Repeat atropine at 5-10 minute intervals until atropinization occurs (dry, flushed skin, tachycardia, pupillary dilatation), and maintain for 48 hours. If cyanotic, give initial atropine intramuscularly and start measures to improve ventilation.

Start 2-PAM (PROTOPAM, Ayerst) at same time. Give 1-2 grams PROTOPAM (20-40 mg/kg for children) in 100cc saline over 15-30 minutes. If pulmonary edema is present, give intravenously slowly as a 5% solution in water over a period of not less than 5 minutes.

A second dose may be given after one hour if muscle weakness persists. Additional doses may be given cautiously for persistent muscle weakness. May be given by intramuscular or subcutaneous routes if intravenous administration is not feasible.

In case of skin contact wash patient with soap and water followed by wash with 95% ethyl alcohol. Keep patient under constant observation for 24-36 hours. Symptoms may persist for one month.

The use of theophylline, morphine, barbiturates, phenothiazines, reserpine, and succinyl choline is contraindicated.

Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests. Repeated exposure to cholinesterase inhibitors may, without warning, cause increasing susceptibility to very small doses of any cholinesterase inhibitor. This will last for several weeks following exposure.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Birds and wildlife feeding in treated areas may be killed. Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not use, pour, spill or store near heat or open flame. Do not store below 32°F.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

CONDITIONS OF SALE, WARRANTY, LIMITATION OF LIABILITY

Rhone-Poulenc warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes described on the label. Rhone-Poulenc makes no other express or implied warranty including any warranty of fitness or merchantability. In no case shall Rhone-Poulenc or the seller be liable for consequential, special or indirect damages such as lost profits or values resulting from the use or handling of this product.

BEST DOCUMENT AVAILABLE