


Product Number
2010



ALLETTE™
FUNGICIDE



ACTIVE INGREDIENT:
Aluminum tris (0-ethyl phosphonate).....80.0%

INERT INGREDIENTS:.....20.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT


IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Rinse with plenty of water. Get medical attention if irritation persists.

See Back Panel For Additional Precautionary Statements.

EPA Reg. No. 359-TNA
EPA Est
-0783

NET CONTENTS
POUNDS



Manufactured by
RHÔNE-POULENC INC.
AGROCHEMICAL DIVISION
Monmouth Junction, N.J. 08852

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands.
Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL:

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL:

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities

GENERAL INFORMATION

NOTE TO USER: During mixing and applying this product, wear full protective clothing (long pants, long sleeve shirt and water resistant gloves).

ORNAMENTALS:

Aliette may be used on field and greenhouse ornamentals.

When used in conjunction with good cultural management practices, Aliette is effective in controlling Phytophthora root rot on the following ornamentals:

Azalea	Japanese Holly
Rhododendron	Monterey Pine
Juniper	Pieris japonica
Pittosporium	Schefflera
Boxwood	

Aliette may be applied either as a drench or by incorporation into the potting mixture. Applications should begin at potting and always prior to disease development. Applications should continue for as long as conditions are favorable for Phytophthora development.

Aliette should be applied on ornamentals in accordance with the table below.

<u>Pest</u>	<u>Rate</u>	<u>Application</u>	<u>Amount</u>	<u>Interval</u>
Phytophthora root rot (<i>Phytophthora</i> sp.)	6.5-13oz/yd ³	Incorporation	-	At Potting
	1-2 lbs/1000 ft ²	Drench	0.5-1.5 pts/ft ²	Monthly

CONDITION OF SALE, WARRANTY, LIMITATION OF LIABILITY

This product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of Rhone-Poulenc or the seller. Buyer assumes all risks of use, storage, or handling of this material, not in strict accordance with directions given herewith. In no case shall Rhone-Poulenc or the seller be liable for consequential, special, or indirect damages such as loss of profits or values resulting from the use or handling of this product.

BEST DOCUMENT AVAILABLE