

| | | |
|---|---------------------------|--|
| U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (PH 567) WASHINGTON, D.C. 20460 | EPA REGISTRATION NO. | DATE OF ISSUANCE NOV 23 1984 |
| | TERM OF ISSUANCE | |
| | NAME OF PESTICIDE PRODUCT | |

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
 (Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered under the Act with the following conditions, provided that you:


1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA Sect. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Place the labeling and/or label on the product in accordance with the Act and Regulations.
3. Add the phrase "EPA Registration No. 509-711".
4. Add - KEEP OUT OF REACH OF CHILDREN, above the statement of hazards, and a reference to side panel precautions - to the front panel of the label.
5. Add the following hazard statement, to the front panel of the label: "Do not spray directly to water." and add the following precautionary statement to the front panel of the label: "Do not spray directly to water."
6. Add the following hazard statement, to the front panel of the label: "Do not spray directly to water."

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL **NOV 26 1984**

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Richard F. Mountford 
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

Product Number
2146

CHIPCO® RONSTAR® 50 WP Herbicide



Selective Preemergent Herbicide for the Control of Annual Grasses
and Broadleaf Weeds in Turf

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND
LANDSCAPE PERSONNEL ONLY. NOT FOR USE ON HOME LAWNS.

ACCEPTED
with comment
NOV 26 1984
The Federal Insecticide,
Fungicide, and Rodenticide
Act
EPA Reg. No. 359-714

ACTIVE INGREDIENT:
Oxadiazon (2-tert-butyl-4-(2,4-dichloro-5-
isopropoxyphenyl)- Δ^2 -1,3,4-oxadiazolin-5-one)..... 50%
INERT INGREDIENTS..... 50%

Keep Out of The Reach of Children

WARNING

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in Eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements

EPA Reg. No. 359-XXX
EPA Est.
1084

NET CONTENTS
POUNDS

RHÔNE-POULENC INC.
AGROCHEMICAL DIVISION
MONTPELLIER, FRANCE 34100

ADD:

Wear protective clothing including long sleeve shirt and gloves when handling. *duplicate*

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Causes substantial but temporary eye injury. Wear goggles or face shield when handling material. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not discharge into lakes, streams or public waters unless in accordance with an NPDES permit. For guidance contact your Regional Office of the U.S. Environmental Protection Agency.

*Do not apply directly to water
Do not contaminate water by cleaning of equipment or disposal of wastes*

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not store near other pesticides or seeds.

PESTICIDE DISPOSAL: *OK RTR*

Wastes resulting from the use of this product may be disposed of on site or at an approved waster disposal facility.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

*Tox Category II
Statement*

GENERAL INFORMATION

Chipco Ronstar 50 WP is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf. Chipco Ronstar 50 WP controls weeds by killing the young seedlings as they come in contact with the herbicide during germination.

Apply Chipco Ronstar 50 WP with properly calibrated application equipment that will apply the spray solution uniformly (uniform application is essential for satisfactory weed control). Calibrate the application equipment prior to use, according to the manufacturer's directions and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid con-

tamination of water intended for irrigation and domestic use.

TURF

For use in established perennial Bluegrass, Bermudagrass, Perennial Ryegrass, St. Augustinegrass and Tall Fescue turf such as fairways, parks, golf courses and lawns. Chipco Ronstar 50 WP controls Crabgrass (*Digitaria* spp), Annual Bluegrass (*Poa annua*), and Goosegrass (*Eleusine indica*); also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis* spp.), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus officinis*) and Pigweed (*Amaranthus* spp.).

Rake turf area free of leaves, mowing if necessary so the spray solution reaches the soil surface. Measure the area to be treated and uniformly apply the amount required. If rain is not expected shortly after application, sprinkle the turf thoroughly.

AMOUNT TO USE

Crabgrass and Goosegrass Control: For preemergent control, apply 4-8 pounds of Chipco Ronstar 50 WP per acre in 30-150 gallons of water or 1.5-3 ounces of Chipco Ronstar 50 WP per 1000 sq.ft. in .75-3.5 gallons of water. Application should be made in early spring prior to seed germination. Application made after March or April may not be fully effective.

Annual Bluegrass Control: For preemergent control, apply 8 pounds of Chipco Ronstar 50 WP per acre in 30-150 gallons of water or 3 ounces of Chipco Ronstar 50 WP per 1000 sq.ft. in .75-3.5 gallons of water in late summer to early fall or in early spring prior to seed germination.

Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass Control: For preemergent control of these weeds, apply 8 pounds of Chipco Ronstar 50 WP per acre in 30-150 gallons of water or 3 ounces of Chipco Ronstar 50 WP per 1000 sq. ft. in .75-3.5 gallons of water in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 6 pounds of Chipco Ronstar 50 WP per acre.

Chipco Ronstar 50 WP may cause discoloration of Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown withing 2 to 3 weeks.

Do not apply Chipco Ronstar 50 WP to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Chipco Ronstar 50 WP should not be used on putting greens or tees because of the varying cultural practices used.

CONDITION OF SALE, WARRANTY, LIMITATION OF LIABILITY

This product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of Rhone-Poulenc or the seller. Buyer assumes all risks of use, storage, or handling of this material, not in strict accordance with directions given herewith. In no case shall Rhone-Poulenc or the seller be liable for consequential, special, or indirect damages such as loss of profits or values resulting from the use or handling of this product.