

105?
1025
116


PRODUCT NUMBER
2073

SUPPLEMENTAL LABELING

BUCTRIL R 4EC HERBICIDE

WEED CONTROL IN COVER CROPS
ON CONSERVATION RESERVE ACRES

DIRECTIONS FOR USE



CROPS:

Crested wheatgrasses (all), tall wheatgrass, western wheatgrass, bluebunch wheatgrass, intermediate wheatgrass, perennial ryegrass, fescues, bentgrasses, Kentucky bluegrass, Canby bluegrass, bahiagrass, Russian wildrye, seedling alfalfa.

TARGET WEEDS:

Buctril 4EC will control broadleaf weeds mentioned in the GENERAL WEED LIST found on the package label. Follow all use directions on this label and on the Buctril 4EC package label to ensure product performance and crop safety.

WHEN TO APPLY:

Buctril 4EC may be applied with standard ground boom and aerial hydraulic spray systems to forage crops listed here anytime after the 2-3 leaf stage, but before weeds exceed recommended stages of growth for treatment. Seedling alfalfa may be treated anytime after the 3-trifoliate leaf stage.

HOW TO APPLY:

Ground Application: Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with flat fan nozzles spaced a maximum of 20 inches apart and screens no finer than 50 mesh in the nozzle tips and in-line strainers. Apply in a minimum spray volume of 10 gallons per acre with minimum spray pressure of 30 psi at the nozzle tips.

Aerial Application: The sprayer should be equipped to provide a uniform and accurate application. Use a minimum spray volume of 5 gallons per acre using diaphragm-type nozzles that produce cone or fan spray patterns with a maximum spray pressure of 40 psi at the nozzle tips. Nozzles should be located no further than 3/4 the distance from the center of the aircraft to the end of the wing or rotor. Nozzles should be oriented at an angle between straight down and straight back. The aircraft should discharge the spray a maximum of 10 feet above the crop. To minimize drift, do not apply during periods of gusty winds or when wind exceeds 5 MPH.

In order to ensure maximum safety and weed control, follow label recommendations on this label and all the cautions and limitations on the package labels of the tank mixture components.

USE RECOMMENDATIONS:

BUCTRIL 4EC

TIMING OF APPLICATION		
RATE	CROP	WEED ^{1/}
3/4 Pint/A	Apply anytime in the fall or spring after grasses have reached the 2-3 leaf stage or when alfalfa has reached the 3-trifoliate leaf stage.	Most Susceptible Broadleaf Weeds. Apply to weeds up to the 8 leaf stage or 4 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
1 Pint/A		Susceptible Broadleaf Weeds. Apply to weeds up to the 4 leaf stage or 2 inches in height whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.

^{1/} Refer to the general weed list on the Bucril 4EC package label for a complete list of susceptible weeds.

BUCTRIL 4EC AND MCPA - Grass Crops Only - Do not use on alfalfa.

A tank mixture of Bucril 4EC at the recommended rate + MCPA at 1/4 to 1/2 pound of active ingredient per acre is used to broaden the spectrum of broadleaf weed control. The tank mixture will control the weeds listed on the MCPA label. In addition, to the weeds listed on the Bucril 4EC label. The rate of Bucril 4EC in the tank mixture is the same as when Bucril 4EC is used alone. See BUCTRIL 4EC RECOMMENDATIONS.

Bucril 4EC + MCPA tank mixtures can be applied after the 3 leaf stage but before the crop reaches the boot stage. Apply only to the following grass species: Crested wheatgrasses, tall wheatgrass, blue bunch wheatgrass, intermediate wheatgrass, sheep fescue, canby bluegrass, Russian wildrye. Do not apply to alfalfa.

USE RESTRICTIONS:

Do not allow livestock to graze treated fields.

Do not feed treated grasses or alfalfa to livestock.

Do not apply Bucril 4EC + MCPA tank-mixtures to alfalfa.

Mix with clean water only, do not use spray additives or fluid fertilizers as crop injury may occur.

Do not apply Bucril 4EC to alfalfa when air temperatures exceed 80°F as crop injury may occur.

This labeling must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, and precautions on the Bucril 4EC and MCPA package labels are to be followed.

