

PRODUCT NUMBER
5030

REPLY AVAILABLE COPY

SPECIMEN LABEL

ACCEPTED
MAY 11 1987
Under Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 359-717

MOCAP[®] - PCNB NEMATODE INSECTICIDE



3-10
GRANULAR

Pre-planting treatment for nematodes, corn rootworm and white mold control on peanuts.

ACTIVE INGREDIENT: Ethoprop	
O-Ethyl S, S-Dipropyl Phosphorodithioate	3%
Pentachloronitrobenzene	10%
INERT INGREDIENTS	87%
TOTAL	100%

U.S. Patent Nos. 3,112,244; 3,268,393

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

ON SKIN: Immediately remove all contaminated clothing. Wash skin with soap and water. Wash Clothing before reuse. Discard contaminated shoes.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to and unconscious person.

IN EYES: Poisonous if in eyes. Flush with plenty of water for at least 15 minutes. Call a physician immediately.

IF INHALED: Remove to fresh air and apply artificial respiration if indicated.

See back Panel for additional Precautionary Statements and Note to Physician

EPA Reg.No. 359-TRT
EPA Est.No.
0486

Manufactured by

RHÔNE-POULENC INC.

Net W
Pc
(

3-4-77

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be Fatal if absorbed through skin. Rapidly absorbed through skin. May cause skin irritation. This product is a possible skin sensitizer. If an allergic reaction occurs after repeated exposure discontinue use and consult a physician.

Harmful if swallowed or inhaled. Do not get in eyes, on skin or on clothing. Wear rubber gloves, long sleeved shirt, long pants, socks and shoes. In case of contact, wash immediately with water.

Do not breathe fumes or dust. After use wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not contaminate feed or food. Do not allow livestock to feed in treated areas.

SYMPTOMS OF POISONING: A sense of "tightness" in chest, sweating, stomach pains, vomiting, diarrhea, contraction of pupils.

NOTE TO PHYSICIAN: This product is a cholinesterase inhibitor. Warning symptoms include weakness, headaches, tightness of chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

Administer atropine, 2 mg, intramuscularly or intravenously. Repeat dose every 3 to 8 minutes until signs of atropinization occur. Give pralidoxime chloride (Protopam or 2-PAM), 2.5 gm in 100 cc of sterile water, intravenously, slowly in 15 to 30 minutes or by deep intramuscular injection of 1 gram in 3 cc of distilled water. Repeat every half hour if respiration weakens or muscle fasciculation or convulsion recur. Clear chest by postural drainage. Artificial respiration may be necessary. Observe patient continuously for 48 hours. Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests. Repeated exposure to cholinesterase inhibitors may, without warning, cause increasing susceptibility to very small doses of any cholinesterase inhibitor.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Birds feeding in treated areas may be killed. Cover or incorporate granules that are spilled. Do not apply directly to water or wetlands. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Unprotected persons may enter treated fields after chemical has been mixed into the soil.

25-8-27

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not store in or around the home.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PEANUT

Row treatment at early pegging for control of nematodes, corn rootworm and white mold (Southern stem rot): Use 100 pounds of MOCAP-PCNB per acre at early pegging. Apply in a band 12 to 15 inches wide centered on the row (36-inch row spacing), or 2/3 pound per 100 linear feet of row. Do not apply when plants are wet. Incorporate lightly into the soil.

NOTE: Do not apply more than 10 pounds of pentachloronitrobenzene (PCNB) in any form in any one season. Do not use treated vines as feed or livestock. Do not hog down peanuts. Do not plant any root crop in rotation on PCNB-treated soil except potatoes.

CONDITION OF SALE, WARRANTY, LIMITATION OF LIABILITY

This product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of Rhone-Poulenc or the seller. Buyer assumes all risks of use, storage, or handling of this material, not in strict accordance with directions given herewith. In no case shall Rhone-Poulenc or the seller be liable for consequential, special, or indirect damages such as loss of profits or values resulting from the use or handling of this product.

3-4-87