

10F4
Pm 21
359-725

ACCEPTED
AUG 11 1987
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 359-725

ROVRAL® 30
FLOWABLE FUNGICIDE

ACTIVE INGREDIENT:

Iprodione; 3-(3,5-dichlorophenyl)-N-
(1-methylethyl)-2,4-dioxo-1-
imidazolidinecarboxamide* - - - - - 30%

INERT INGREDIENTS - - - - - 70%

Equivalent to 2.79 lbs. active Iprodione per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT
In case of contact, wash skin with soap and
water; for eyes flush with water for 15 minutes
and get medical attention.
See back panel for additional precautionary statements

EPA Reg. No. 359
EPA Est. 7501-1-1

NET CONTENTS
Gallons
Liters

Manufactured by

RHONE-POULENC INC.
Agrochemical Division
Monmouth Junction, New Jersey 08852

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

C A U T I O N
Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. *And copy A*

ENVIRONMENTAL HAZARDS
Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTION FOR USE
GENERAL USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Add copy B →

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.


Rovral 30 Flowable Fungicide, when used as directed, protects turfgrass and ornamental seed and seedlings from seed-borne and soil-borne fungi causing seed decay, damping-off, and seedling blights. Rovral 30 is particularly effective against Alternaria, Botrytis, Fusarium, Helminthosporium, Rhizoctonia, and Sclerotinia.

For additional control of Pythium damping-off, it is recommended that Rovral 30 be applied in combination with Gustafson APRON® FL in accordance with labeled uses on turfgrass.

Seeds should be well cured and cleaned prior to treatment. Use only the recommended rates of product per 100 pounds of seed.


<u>Planting Seeds</u>	<u>Application Rates per 100 lbs. of Seed</u>	<u>Seedling Disease Complex</u>
Turfgrass	3.0 to 6.0 Fl. Oz. (88.8 to 177.6 cc)	<u>Fusarium nivale</u> , <u>Fusarium rosei</u> , <u>Helminthosporium spp.</u> , <u>Rhizoctonia solani</u> , and <u>Sclerotinia hemoecarpa</u>
Ornamentals	3.0 to 6.0 fl. oz. (88.8 to 177.6 cc)	<u>Alternaria spp.</u> , <u>Ascochyta</u> <u>chrysanthemi</u> , <u>Botrytis spp.</u> , <u>Rhizoctonia spp.</u>

Thoroughly mix Rovral 30 Flowable Fungicide into the required amount of water for the slurry treater equipment and dilution rate used. Clean equipment thoroughly between treating operations.



Important: Before using - stir thoroughly to mix contents or use entire container at one time. Do not combine colloidal suspensions, wettable powders, or emulsifiable concentrates unless previous experience assures compatibility.

Note: Although no phytotoxicity has been observed on seeds or seedlings treated at recommended use rates, it is recommended that evaluations be conducted on the particular species and varieties of seed to be treated prior to commercial application.



Copy A

Avoid breathing vapors or spray mist.

Copy B

Do not use treated seed for food, feed, or oil purposes.

Since this product does not contain a dye, all seed treated with this product (except by the hopper box method) must be colored with an EPA-approved dye (e.g. 40 CFR 180.1001) which imparts an unnatural color to the seed to help prevent the inadvertent use of treated seed for food, feed, or oil purposes.

The Federal Government and most states have seed laws or regulations requiring that certain caution statements be affixed to containers used to ship or store treated seed.

