

NET WEIGHT 1 OZ.

side panel.

## DIRECTIONS

Place teaspoon quantities of bait at 8 to 10 foot intervals where mice feed, water and/or travel. Place bait in bait boxes constructed of wood or metal, tamperproof and with baffled entrances. Remove all other food sources. Collect and burn or bury all uneaten baits after 72 hours. Collect and burn or bury dead animals. Do not treat the same area with strychnine in less than 60-day intervals. Acceptable bait materials are hulled brains, oats, barley, wheat, coated with 0.3%to 0.5% active ingredient. Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamperproof bait boxes. Poisonous if swallowed. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Dispose of excess treated seeds or wastes by burial. Apply this product only as specified on this label. DANGER: CONVULSIVE FOISON. Do not contaminate feed and Foodstuffs. Keep away from children and domestic animals.

ANTIDOTE: If less than 10 minutes has passed since the poison was taken, give a tablespoonful of salt in a glass of warm water. Have victim lie down in a quiet darkened room and keep him warm.

CALL A PRYSICIAN IMMEDIATELY



POISON	Active Ingredients:
(Skull and crossbones)	Strychnine Alkaloid 0.5%
	Inert Ingredients
	Total $100.0\%$

E.P.A. Reg. No. 368-30