

★ STAR

Star spells RATS backwards

CONTAINS PROLIN[★]

READY TO USE • MONEY BACK GUARANTEE

Kills RATS and MICE

**CAUTION: Keep out of the reach of children.
See other caution on side panel.**

PELLET FORM

E.P.A. No. 368-39

1 lb. \$1.00; 2 1/2 lb. \$1.69; 5 lb. \$2.98 50 lb. \$27.50; 100 lb. \$50.00

ACTIVE INGREDIENTS:

Warfarin (3[ALPHA-ACETONYLBENZYL]-4-HYDROXYCOUMARIN)	0.025 %
N1-(2-QUINCOXALINYL) SULFAMLAIDE(SULFAQUINOXALINE)	0.025 %
Inert ingredients	99.950 %
Total	100.000 %

*Prolin is a registered trademark of the Wisconsin Alumni Research Foundation for a rodenticide.

GET THE ECONOMICAL 5lb. SIZE

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

DO NOT BE USED IN AREAS INACCESSIBLE TO CHILDREN, WILDLIFE AND DOMESTIC ANIMALS OR IN TAMPER-RESISTANT BAIT.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

If swallowed by humans, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intra-venous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bis-hydroxycoumarin. In case of ingestion, call a physician.

MASTER LABORATORIES, BEAVER FALLS, PA. E.P.A. NO. 368-39

INSTRUCTIONS

RATS: Place 4 to 6 ounces of bait in dry container next to burrows, runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 15 days and continue baiting until all signs of feeding have stopped. Replace stale bait, damp or moldy baits with fresh bait.

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feeding, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin. Achieve better results with proper baiting. Place baits along runways, behind boards, against wall. Keep bait dry. Open bugs and use as bait stations.

468-39

Str

CCF

READY TO

Rat

CAUTION:
See other

1 lb. \$1.00; 2 1/2

ACTIVE INGREDIENTS
Warfarin (3) (AL)
N1-(2-QUINCO
Inert Ingredients

*Prolin is a registered trademark for a rodenticide

GET TH

MASTER

RED

RED

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

BAITS SHOULD BE PLACED IN AREAS INACCESSIBLE TO CHILDREN, PETS WILDLIFE AND DOMESTIC ANIMALS OR IN TAMPER PROOF BAIT BOXES.

RED

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

If swallowed by humans, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intra-venous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin. In case of ingestion, call a physician.

MASTER LABORATORIES, BEAVER FALLS, PA. E.P.A. NO. 368-39

RED

INSTRUCTIONS

RATS: Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 15 days and continue baiting until all signs of feeding have stopped. Replace stale bait, damp or moldy baits with fresh bait.

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin.

Achieve better results with proper baiting. Place baits along runways, behind boards, against wall. Keep bait dry. Open bags and use as bait stations.

ACCEPTED
JAN 1 1974
468-39

RED