

warning!

MAY BE FATAL IF SWALLOWED

Avoid breathing dust.

Do not get in eyes or on skin.

Wash thoroughly after handling.

Discard container immediately when empty. Do not reuse.

Do not graze treated areas. Do not feed clippings to livestock.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

EPA Registration No. 372-24

5648

Mallinckrodt

25 LBS.

(11.3 Kgs.)

kromad

ACTIVE INGREDIENTS

Cadmium Sebacate	5	%
Potassium Chromate	5	%
Malachite Green	1	%
Auramine	0.5	%
Thiram (Tetramethylthiuram Disulfide)	16	%
Total Cadmium - Cd	1.6	%
Total Chromium - Cr	1.3	%

INERT INGREDIENTS

72.5 %
Urea • Ferrous Sulfate • Wetting Agent

WARNING!

KEEP OUT OF REACH OF CHILDREN.

See side panels for additional cautions.

Manufactured for

MALLINCKRODT CHEMICAL WORKS

ST. LOUIS, MO. 63160

ACCEPTED

DECEMBER 11, 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION UNDER NO. 372-24

KROMAD

KROMAD prevents and controls brown patch, dollar spot, copper spot, red thread (pink patch), and leaf blights, such as incited by Helminthosporium and Curvularia. KROMAD contains a fungicidal dye to assist in disease control and to provide turf color.

Comments

directions

APPLICATION RATE

PREVENTIVE — Under normal weather conditions, for the prevention of disease, apply KROMAD every 5 to 7 days at 3 ounces per 1000 square feet. During weather particularly favorable for disease, such as high temperature and humidity, apply KROMAD more frequently or at heavier rates. It may be applied as often as 3 or 4 times per week, or at rates of 4, 6 or even 8 ounces per 1000 square feet, as necessary to keep disease under control, without injury to fine turf grasses.

CURATIVE — If disease is already active, apply KROMAD at 5 or 6 ounces per 1000 square feet every 3 to 7 days until disease is under control. Then follow preventive schedule.

SPRAY PREPARATION

Fill tank with five to ten gallons of water for each 1,000 sq. ft. of turf to be treated. Add required amount of KROMAD and, where possible, keep in suspension by agitation.

IMPORTANT—Do not water in after spraying. Fungicides have a tendency to wash from grass blades during hard rains or under conditions of heavy daily watering. Watering after application is not necessary and preferably should be held off for 24 hours. If there is danger that KROMAD has been removed, it should be re-applied, especially during periods when diseases are expected to be severe.

U.S. Pat. 2,876,155


Printed in U.S.A. 107218-1

ACCEPTED WITH COMMENTS

5648



25 LBS. (11.3 Kgs.)



kromad®

ACTIVE INGREDIENTS

- Cadmium Sulfate
- Phosphoric Acid
- Magnesium
- Azoxystrobin
- Triphenyltin Hydroxide
- Iron Oxide

INERT INGREDIENTS

- Iron Oxide
- Ferrous Sulfate
- Wetting Agent

WARNING!

KEEP OUT OF REACH OF CHILDREN.

See label panels for additional cautions.

Manufactured for

MALLINCKRODT CHEMICAL WORKS

ST. LOUIS, MO. 63160

MAY BE FATAL IF SWALLOWED

Avoid breathing dust.

Do not get in eyes or on skin.


Wash thoroughly after handling.

Discard container immediately when empty. Do not reuse.

Do not graze treated areas. Do not feed clippings to livestock.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

EPA Registration No. 372-24



KROMAD

KROMAD prevents and controls brown patch, dollar spot, copper spot, red thread (pink patch), and leaf blights, such as incited by Helminthosporium and Curvularia. KROMAD contains a fungicidal dye to assist in disease control and to provide turf color.

APPLICATION RATE

PREVENTIVE: In preventive applications, weather conditions are favorable for disease. Apply KROMAD every 10 to 14 days at 5 ounces per 1000 square feet. During weather conditions favorable for disease, such as high temperature and humidity, apply KROMAD more frequently at 1 to 2 ounces per 1000 square feet. To prevent diseases per 1000 square feet, apply KROMAD to keep disease under control. A minimum of 1 to 2 ounces per 1000 square feet is recommended.

CURATIVE: In curative applications, apply KROMAD at 1 to 2 ounces per 1000 square feet every 10 to 14 days until disease is under control. A minimum of 1 to 2 ounces per 1000 square feet is recommended.

SPRAY PREPARATION

Fill tank with five to ten gallons of water for each 1000 sq. ft. of turf to be treated. Add required amount of KROMAD and, where possible, keep in suspension by agitation.

IMPORTANT: Do not water in after spraying. Fungicides have a tendency to wash from grass blades during hard rains or under conditions of heavy daily watering. Watering after application is not necessary and preferably should be held off for 24 hours. If there is danger that KROMAD has been removed, it should be re-applied, especially during periods when diseases are expected to be severe.

U.S. Pat. 2,876,155

Printed in U.S.A. 107216 1