

Warning!

MAY BE FATAL IF SWALLOWED

Avoid breathing dust.

Do not get in eyes or on skin.

Wash thoroughly after handling.

Discard container immediately when empty. Do not reuse.

Do not graze treated areas. Do not feed clippings to livestock.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Mallinckrodt
MALLINCKRODT, INC.

5648

Mallinckrodt

25 LB. (11.3 Kg)



ACTIVE INGREDIENTS

Calcium Sulfate
Potassium Chloride
Triphenyl Tinylate
Total Fungicide

INERT INGREDIENTS


Urea • Ferrrous Sulfate • Wetting Agent

WARNING!

KEEP OUT OF REACH OF CHILDREN

EPA Registration No. 372-24

EPA Establishment No. 31429 USA



KROMAD

KROMAD prevents and controls brown patch, dollar spot, copper spot, red thread, pink patch, and leaf blights such as incited by Helminthosporium and Sclerotinia. KROMAD contains a pigment to aid uniform application and to provide turf color.

direction

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

FOR PROFESSIONAL USE ONLY

APPLICATION RATE

PREVENTIVE — Under normal weather conditions for the prevention of disease, apply KROMAD every 5 to 7 days at 3 oz. per 1000 sq. ft. During weather particularly favorable for disease such as high temperature and humidity, apply KROMAD more frequently or at heavier rates. It may be applied as often as 3 or 4 times per week, or at rates of 4, 6 or even 8 oz. per 1000 sq. ft. as necessary to keep disease under control, without injury to fine turf grasses.

CURATIVE — If disease is already active, apply KROMAD at 5 or 6 oz. per 1000 sq. ft. every 3 to 7 days until disease is under control. Then follow preventive schedule.

SPRAY PREPARATION

Fill tank with five to ten gal. of water per sq. ft. of turf to be treated. Add required amount of KROMAD and where possible keep in agitation.

IMPORTANT — Do not water in areas where fungicides have a tendency to wash blades during hard rains or under conditions of heavy watering. Watering after application is necessary and preferably should be done during the day. If there is danger that KROMAD will be washed away, it should be re-applied, especially during periods when diseases are expected to recur.

WARRANTY

This material conforms to the description on the label and is reasonably fit for the purposes intended. The effectiveness of use, timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, compatibility with other chemicals, and other factors are beyond the control of the manufacturer. Buyer assumes all responsibility for the use of this material. No warranty is made by the manufacturer for the use of this material in accordance with directions on the label. **OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS DENIED.**

5648

Mallinckrodt

25 LB. (11.3 Kg)

KROMAD

ACCEPTED

JUN 1 1980

KROMAD prevents and controls brown patch, dollar spot, copper spot, red thread, pink patch, and leaf blights, such as caused by Helminthosporium and Curvularia. KROMAD contains a pigment to aid uniform application and to provide turf color.

kromad

BROAD SPECTRUM TURF FUNGICIDE

ACTIVE INGREDIENTS

Cadmium Sebacate
Potassium Chlorate
Thiram
Total Cadmium
Total Chlorine

INERT INGREDIENTS

Urea • Ferric Sulfate • Wetting Agent

WARNING!

KEEP OUT OF REACH OF CHILDREN

EPA Registration No. 372-24

EPA Establishment No. 32429-GA-1

directions

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

FOR PROFESSIONAL USE ONLY

APPLICATION RATE

PREVENTIVE - Under normal weather conditions for the prevention of disease, apply KROMAD every 5 to 7 days at 3 oz. per 1000 sq. ft. During weather particularly favorable for disease, such as high temperature and humidity, apply KROMAD more frequently or at heavier rates. It may be applied as often as 3 or 4 times per week, or at rates of 4, 6 or even 8 oz. per 1000 sq. ft. as necessary to keep disease under control without injury to fine turf grasses.

CURATIVE - If disease is already active, apply KROMAD at 5 or 6 oz. per 1000 sq. ft. every 3 to 7 days until disease is under control, then follow preventive schedule.

SPRAY PREPARATION

Fill tank with five to ten gal. of water for each 1000 sq. ft. of turf to be treated. Add required amount of KROMAD and, where possible, keep in suspension by agitation.

IMPORTANT - Do not water in after spraying. Fungicides have a tendency to wash from grass blades during hard rains or under conditions of heavy daily watering. Watering after application is not necessary and preferably should be delayed for 24 hours. If there is danger that KROMAD has been removed, it should be re-applied, especially during periods when diseases are expected to be severe.

WARRANTY

The material herein is the result of the best knowledge and skill of the manufacturer at the time of application. The manufacturer does not warrant the results of the application of this material under any conditions. The user assumes all liability for the results of the application of this material. The manufacturer does not warrant the results of the application of this material under any conditions. The user assumes all liability for the results of the application of this material.

124