

AGWAY PROTECTOR-3L SEED TREATMENT

ACTIVE INGREDIENT

2-(thiocyanomethylthio) benzothiazole

INERT INGREDIENTS

30.0%

70.0%

100.0%

DANGER: KEEP OUT OF REACH OF CHILDREN.

See additional precautionary statements on side panel

GENERAL INFORMATION

Protector is recommended for the control of soil and seed borne fungi which impair the germination and seedling development of field corn, safflower, sorghum, cotton, sugar beet, oats, rice and wheat. Protector when used as directed will usually increase stands and yields of field corn, oats, rice and wheat by preventing seed decay and seedling diseases.

DIRECTIONS:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage, cleaning of equipment or disposal. Open dumping is prohibited.

Pesticide, spray mixture or rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place.

Consult federal, state or local disposal authorities for approved alternative procedures.

Field Corn Seed Protection For ready mix treaters use 2.0 fluid ounces of Protector per 100 pounds of corn seed. Protector may be dispersed in water for distribution provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters disperse 2.0 fluid ounces of Protector in 8 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater provided that 2.0 fluid ounce of Protector is applied per 100 pounds of seed.

Safflower Seed Protection For ready mix treaters use Protector 3L at the rate 1.0 to 3.0 fl. oz. per 100 lb. of seed. Protector 3L may be dispersed in water for distribution provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters disperse 1.0 to 3.0 fl. oz. of Protector 3L in 8 fl. oz. of water and apply this dispersion per 100 lb. of seed.

However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating provided that Protector 3L is applied at 1.0 to 2.0 fl. oz. per 100 lb. of seed.

Sorghum Seed Protection For ready mix treaters use 2.5 fl. oz. per 100 lb. of seed. Protector 3L may be dispersed in water for distribution provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters disperse 2.5 fl. oz. of Protector 3L in 8 fl. oz. of water and apply this dispersion per 100 lb. of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating provided that 2.5 fl. oz. of Protector 3L is applied per 100 lb. of seed.

Sugarbeet Seed Protection For ready mix treaters use Protector 3L at the rate of 3.0 to 4.0 fl. oz. per 100 lb. of seed. Protector 3L may be dispersed in water for distribution provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters disperse 3.0 to 4.0 fl. oz. of Protector 3L in 8 fl. oz. of water and apply this dispersion per 100 lb. of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating provided that Protector 3L is applied at 3.0 to 4.0 fl. oz. per 100 lb. of seed.

Cotton Seed Protection For ready mix treaters use Protector 3L undiluted at the rate of 4.0 to 5.0 fl. oz. per 100 lb. of acid delinted seed and 5.0 to 6.0 fl. oz. per 100 lb. of machine delinted seed. Protector 3L may be diluted with water for distribution provided the recommended amount of product is applied. For slurry treater prepare slurry by stirring 1.6 to 2.0 pints (for acid delinted seed) of Protector 3L in water to a final volume of 1 gallon. Suggested use is 0.125 pint of the slurry per 10 lb. of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating provided that Protector 3L is applied at 4.0 to 5.0 fl. oz. per 100 lb. of acid delinted seed or 5.0 to 6.0 fl. oz. per 100 lb. of machine delinted seed.

Barley, Oats, Rice and Wheat Seed Protection For ready mix treaters use 0.75 fluid ounce of Protector per bushel of seed. Protector may be dispersed in water for distribution provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters disperse 0.75 fluid ounce of Protector in 4 fluid ounces of water and apply this dispersion per bushel of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater provided that 0.75 fluid ounce of Protector is applied per bushel of seed.

DISCLAIMER Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOTE This pesticide is to be sold **ONLY** in this original unbroken package.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. If swallowed, call a physician immediately, give patient doses of powdered charcoal immediately or all he can swallow of raw egg whites, milk, gelatin, or flour and water. Then induce vomiting with soap or mustard in warm water. May be harmful if absorbed through the skin. Workmen handling the product should wear rubber gloves and goggles and should avoid contact of the product with clothing or skin. After exposure, wash promptly and thoroughly with an abrasive soap and lots of water and finally with glycerin. In case of persistent irritation of the skin, obtain medical attention. If the product gets into the eyes, it causes eye damage; therefore, flush eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any irritation persists. Keep the drum closed when not in use.

Do not use crops grown from Protector 3L treated seed for grazing of live stock or for harvest as silage or forage for seven weeks after planting. However, grain harvested from plants grown from Protector treated seed can be used for food for feed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water.

Label treated seed. Seeds treated with Protector should not be used for food, feed, and production of any other purpose except planting.

Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 8590-424
AGWAY INC., CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

EPA Est. No. B-000001

A-5-80

NET CONTENTS: 5 GALLONS

70