

4645

Mallinckrodt

25 LBS.

(11.3 Kgs.)

w a r n i n g !


**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
MAY CAUSE SKIN IRRITATION**

Avoid breathing dust or spray mist.
Avoid contact with skin.
Wash thoroughly after handling.
Repeated contact may be harmful.
Do not contaminate or store near feed or foodstuffs.
Do not feed clippings from treated areas to animals.
Do not allow animals to graze on treated areas.

Licensed under U.S. Patents
3,260,588 and 3,260,275

EPA Registration No. 372-34

ACCEPTED
10-2-72
FEDERAL INSTITUTE
FOR ENVIRONMENTAL
HEALTH SAFETY
372-34
SUBJECT
TO COMMENTS



ACTIVE INGREDIENT
5-Ethoxy-3-Trichloromethyl-1,2,4-Thiadiazole.....35%
INERT INGREDIENTS.....65%

WARNING!

KEEP OUT OF REACH OF CHILDREN

See left panel for additional cautions.

MALLINCKRODT CHEMICAL WORKS
ST. LOUIS, MO. 63160

PRODUCT INFORMATION

KOBAN is intended for use on ornamental turf which is subject to Pythium blight, damping off or cottony blight. It may be used at time of seeding to control damping off, or on established turf to prevent or control Pythium blight (cottony blight).

KOBAN does not afford broad spectrum fungus disease control for Rhizoctonia (brown patch), Sclerotinia (dollar spot), Gloeocercospora (copper spot), and similar turf diseases. A fungicide program to control these diseases should be followed if they co-exist with Pythium blight.

DIRECTIONS FOR USE:

For control of Pythium blight (cottony blight, grease spot, damping off) on turf:

Add KOBAN to the spray tank with the agitator in operation. Disperse 4 to 6 ozs. of KOBAN in 5 gallons of water for each 1000 sq. ft. of turf to be treated. Apply uniformly to the turf area with a SPRAY HAWK mobile spray boom or other suitable equipment.

Re-treat in 5-10 days, depending upon climatic conditions, to maintain disease control. KOBAN should not be mixed with other pesticides or nutrients.

For control of damping-off in newly seeded areas: Apply 6-8 ozs. of KOBAN in 5 to 10 gallons of water/1000 sq. ft. immediately after seeding. Irrigate promptly for improved soil penetration of the fungicide and to initiate seed germination. Re-treat at 5 to 7 day intervals at the 4 to 6 oz. rate in order to maintain a preventive disease control program after germination.

Ryegrass Seed Treatment: (Dust) To prevent Pythium Blight add 4 to 6 ozs. KOBAN to the seed treater for each 100 lbs. of ryegrass seed and allow to blend until the KOBAN is thoroughly dispersed in the seed.

IMPORTANT: Bags containing treated seed should bear a tag. This seed has been treated with 35% Ethazole at the rate of 4 to 6 ozs./100 lbs. seed. **DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES.**

Printed in U.S.A. 77231-1

ACCEPTED WITH COMMENTS

Specimen

372-34

4645

Mallinckrodt

25 LBS.


(11.3 Kgs.)

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
MAY CAUSE SKIN IRRITATION**

Avoid breathing dust or spray mist.
Avoid contact with skin.
Wash thoroughly after handling.
Repeated contact may be harmful.
Do not contaminate or store near feed or foodstuffs.
Do not feed clippings from treated areas to animals.
Do not allow animals to graze on treated areas.

Licensed under U.S. Patents
3,260,588 and 3,260,275

EPA Registration No. 372-34



ACTIVE INGREDIENT
5-Ethoxy-3-Trichloromethyl-1,2,4-Thiadiazole 35%
INERT INGREDIENTS 65%

WARNING!
KEEP OUT OF REACH OF CHILDREN

See left panel for additional cautions.

MALLINCKRODT CHEMICAL WORKS
ST. LOUIS, MO. 63160

PRODUCT INFORMATION

KOBAN is intended for use on ornamental turf which is subject to Pythium blight damping off or cottony blight. It may be used at time of seeding to control damping off, or on established turf to prevent or control Pythium blight (cottony blight).

KOBAN does not afford broad spectrum fungus disease control for Rhizoctonia (brown patch), Sclerotinia (dollar spot), Gloeocercospora (copper spot), and similar turf diseases. A fungicide program to control these diseases should be followed if they co-exist with Pythium blight.

DIRECTIONS FOR USE

For control of Pythium blight (cottony blight, grease spot, damping off) on turf:

Add KOBAN to the spray tank with the agitator in operation. Disperse 4 to 6 ozs. of KOBAN in 5 gallons of water or each 1000 sq. ft. of turf to be treated. Apply uniformly to the turf area with a SPRAY HAWK mobile spray boom or other suitable equipment.

Re-treat in 5-10 days, depending upon climatic conditions, to maintain disease control. KOBAN should not be mixed with other pesticides or nutrients.

For control of damping-off in newly seeded areas: Apply 6-8 ozs. of KOBAN in 5 to 10 gallons of water 1000 sq. ft. immediately after seeding. Irrigate promptly for improved soil penetration of the fungicide and to initiate seed germination. Re-treat at 5 to 7 day intervals at the 4 to 6 oz. rate in order to maintain a preventive disease control program after germination.

Ryegrass Seed Treatment: (Dust) To prevent Pythium Blight add 4 to 6 ozs. KOBAN to the seed treater for each 100 lbs. of ryegrass seed and allow to blend until the KOBAN is thoroughly dispersed in the seed.

IMPORTANT: Bags containing treated seed should bear a tag. This seed has been treated with 35% Ethazole at the rate of 4 to 5 ozs./100 lbs. seed. **DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES.**

Printed in U.S.A. 772311

10-2-72
373-34

Handwritten scribbles

Faint printed text at the bottom of the page.