


7693

Mallinckrodt

2 LB.  
(0.9 Kg)



**ACTIVE INGREDIENT -**

5-Ethoxy-3-trichloromethyl- 1,2,4,-thiadiazole*	30%
--	-----

<b>INERT INGREDIENTS</b>	<b>70%</b>
--------------------------	------------

**KEEP OUT OF REACH OF CHILDREN**

**WARNING!**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED.** induce vomiting immediately by giving two glasses of water and sticking finger down throat. Never give anything by mouth to an unconscious person. **GET MEDICAL ATTENTION.**

**IF INHALED.** Induce coughing  
**IF ON SKIN.** Wash off with soap and water.  
**IF IN EYES.** Flush eyes with water.

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

\*Licensed under U.S. Patents 3,260,588 & 3,230,725

Manufactured For

**Mallinckrodt, Inc.**  
 St. Louis, Missouri 63147.

**EPA Reg. No. 372-34**

**EPA Est. No. 228-IL-1**

# KOBAN® 30

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING: HARMFUL IF SWALLOWED, inhaled, or absorbed through skin.** Always wear protective clothing and respirator when mixing and applying pesticides. May cause irritation. Avoid breathing spray mist. Avoid contact with skin, eyes and wet clothing. **IN CASE OF CONTACT,** immediately flush eyes or skin with plenty of water. **GET MEDICAL ATTENTION** if irritation persists. Wash thoroughly with soap and water after handling. Wash contaminated clothing before re-use. Do not contaminate feed or food. Keep children and pets out of treated area during application.

### ENVIRONMENTAL HAZARDS

**THIS PRODUCT IS TOXIC TO FISH.** Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

### FOR PROFESSIONAL USE ONLY

KOBAN 30 is intended for use on ornamental turf which is subject to Pythium blight, damping off or cottony blight. It may be used at time of seeding to control damping off, or on established turf to prevent or control Pythium blight (cottony blight).

KOBAN 30 does not afford broad spectrum fungus disease control for Rhizoctonia (brown patch), Sclerotinia (dollar spot), Gloeocercospora (copper spot), and similar turf diseases. A fungicide program to control these diseases should be followed if they co-exist with Pythium blight.

For control of Pythium blight, cottony blight and damping off on turf.

Night temperatures above 65° F (18° C), with high humidity, are favorable for Pythium development.

Add KOBAN 30 to the spray tank with the agitator in operation. Disperse 2 to 4 oz. of KOBAN 30 in 5 gal. of water for each 1000 sq. ft. of turf to be treated. Use 2 oz./1000 sq. ft. when disease incidence is expected to be light and 4 oz./1000 sq. ft. for more severe conditions.

Apply uniformly to the turf area with a SPRAY-HAWK® mobile spray boom or other suitable equipment.

Re-treat in 5-10 days, depending upon climatic conditions, to maintain disease control. KOBAN 30 should not be mixed with other pesticides or nutrients.

For control of damping-off in newly seeded areas. Apply 7 to 9 oz. of KOBAN 30 in 5 to 10 gal. of water/1000 sq. ft. immediately after seeding. Irrigate promptly for improved soil penetration of the fungicide and to initiate seed germination. Re-treat at 5 to 7 day intervals at the 2 to 4 oz. rate in order to maintain a preventive disease control program after germination.

Use FUNGO-50, KROMAD®, DUOSAN® or CADMINATE® to control a broad spectrum of diseases other than Pythium.

**Grass Seed Treatment:** To prevent Pythium blight add 4 to 7 oz. of KOBAN 30 to the seed treater for each 100 lb. of grass seed. Blend until the KOBAN 30 is thoroughly dispersed in the seed.

**NOTE:** For smaller seeds use the higher rate of KOBAN 30.

**IMPORTANT:** Bags containing treated seed should bear a tag with the following statement: This seed has been treated with 30% Ethazole at the rate of 4 to 7 oz./100 lb./seed. **DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES.**

## STORAGE AND DISPOSAL

**STORAGE:** Not sensitive to light. Stable at normal temperatures and humidities. Store in a dry place at temperature above 45°F (7.2°C) in a well closed container.

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL:** Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

## WARRANTY

This material conforms to the description of the same as reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, fertility, medium, the disease problem, condition of the turf, and compatibility with other chemicals not specifically intended, and other pertinent factors in the use of this product are beyond the control of the manufacturer. However, the manufacturer does warrant that the material is a true product according to the directions set forth herein. **UNLIM. WARRANTY - 4-1-75 (S) OR MERCH. TAB., THIS MADE**

Printed in U.S.A. 8-87