



7693

2 LB.
(0.9 Kg)



ACCEPTED
MAR 19 1982
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
this pesticide
is registered under
EPA Reg. No. 372-34

BEST DOCUMENT AVAILABLE

ACTIVE INGREDIENT
5-Ethoxy-3-trichloromethyl-
1,2,4-thiadiazole* 30%
INERT INGREDIENTS 70%

**KEEP OUT OF REACH OF CHILDREN
WARNING!**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, Induce vomiting immediately by giving
two glasses of water and sticking finger down throat.
Never give anything by mouth to an unconscious person.
GET MEDICAL ATTENTION.

IF INHALED, Induce coughing.

IF ON SKIN, Wash off with soap and water.

IF IN EYES, Flush eyes with water.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

*Licensed under U.S. Patents
3,260,588 and 3,260,725

Manufactured for
MALLINCKRODT, INC.
ST. LOUIS, MISSOURI 63147
EPA Registration No. 372-34
EPA Establishment No. 228-IL-1

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: HARMFUL IF SWALLOWED Irritated or absorbed through skin. Always wear protective clothing and respirator when mixing and applying pesticides. May cause irritation. Avoid breathing spray mist. Avoid contact with skin, eyes and on clothing. **IN CASE OF CONTACT**, immediately flush eyes or skin with plenty of water. **GET MEDICAL ATTENTION** if irritation persists. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets out of treated area during application.

ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO FISH. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

FOR PROFESSIONAL USE ONLY

TRUBAN® is a soil fungicide for control of damping off, root and stem rot diseases caused by Pythium and Phytophthora. It may be used at time of seeding and transplanting either as a soil mix or as a drench for control of these problem fungi in greenhouses and outdoors.

TRUBAN suspensions in water should be constantly agitated to assure uniform application of material. TRUBAN suspensions need not be protected from light. Use at recommended rates. Overdosing may result in toxicity to sensitive plants. If plants are suspect of being sensitive, apply fungicide to a small number to insure safety. TRUBAN should not be mixed with other pesticides or nutrients.

Thoroughly rinse TRUBAN residues out of all equipment after use.

PLANT TYPE	REMARKS
Bedding Plants Agapanthus Anemone Aster Begonia Bougainvillea Calceolaria Celosia Coleus Dahlia Dianthus Foxglove	BEDDING PLANTS - 2 to 3 in. Mix 4 to 12 oz. TRUBAN 30-W with 100 gal. of water and apply to mix to 100 sq. ft. of bed area. Use at time of seeding and transplanting. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Retreat at 4 to 12 week intervals if additional applications are necessary.
Foliage Plants Agave Anthurium Brassia Caladium Chinese Evergreen Dieffenbachia Echeveria Nephthytis Peperomia Pilea Pothos Philodendron Zygocactus	BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS peat and other high organic growth media 6" deep. Mix 8-12 oz. with 100 gal. of water and apply to 400 sq. ft. of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Retreat at 2 to 4 week intervals to maintain control.
	BEDGROWN PLANTS Mix 3 to 10 oz. with 100 gal. of water per 400 sq. ft. or apply in sufficient amount to saturate the soil mixture. Retreat at 4 to 12 week intervals if necessary.
Container & Bedgrown HERBACEOUS - African Violet Begonia - Calceolaria Carnation - Chrysanthemum Easter Lily - Geranium Gerbera - Gypsophila Orchids - Sinningia WOODY Arctostaphylos - Camellia Carrissa - Ceanothus - Fir Hemlock - Juniper - Magnolia Poinsettia - Rhamnus Rhododendron - Southern Pine Western Pine	CONTAINER & BEDGROWN PLANTS Mix 3 to 10 oz. with 100 gal. of water per 400 sq. ft. or apply in sufficient amount to saturate the soil mixture, such as 1/2 pint per 6" pot. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Retreat at 4 to 12 week intervals if necessary.
DRY SOIL MIX for Plants Listed above - Mix 1 to 3 oz. of TRUBAN 30-W with each cu. yd. of soil mixture. Ingredients for the soil mix should be free of large lumps and dry enough to prevent agglomeration during mixing. Improved uniformity of blending will be assured if the TRUBAN is pre-mixed with about five times its volume of dry soil. Retreat if needed as indicated under appropriate soil drench recommendations.	

STORAGE AND DISPOSAL

STORAGE Not sensitive to light. Stable at normal temperatures and humidities. Store in a dry place.

PROHIBITIONS Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL Pesticide spray mixture or rinseate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

WARRANTY

This material contains the best information available at the time of publication. It is not intended to constitute a contract. The user assumes all responsibility for the results obtained. The manufacturer is not liable for any damage or loss resulting from the use of this product.