

4324

2 LBS.



TRUBAN^(H)
FUNGICIDE

30% WETTABLE POWDER
 For control of damping-off root and stem rot diseases caused by Pythium and Phytophthora in ornamental and nursery crops.

ACTIVE INGREDIENT
 5-Ethoxy-3-trichloro-methyl-1,2,4-thiadiazole* 30%

INERT INGREDIENTS 70%

9-27-72
372-42

WARNING!

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED OR
ABSORBED THROUGH SKIN
MAY CAUSE SKIN IRRITATION**

Avoid breathing dust or spray mist.
Avoid contact with skin.
Wash thoroughly after handling.
Repeated contact may be harmful.
Do not contaminate or store near feed or foodstuffs.
Do not use on crops to be used for food.
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

EPA Reg No 372-43
*Licensed under U.S. Patents 3,260,588 & 3,260,725
Manufactured For

Mallinckrodt Chemical Works
St. Louis, Missouri 63160

4324

2 LBS.



TRUBAN
FUNGICIDE

30% WETTABLE POWDER
For control of damping off root and stem rot diseases caused by Pythium and Phytophthora in ornamental and nursery crops.

ACTIVE INGREDIENT

5-Ethoxy-3-trichloro-methyl-1,2,4-triazole*	30%
Inert Ingredients	70%

WARNING!

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED OR
ABSORBED THROUGH SKIN
MAY CAUSE SKIN IRRITATION



Avoid breathing dust or spray mist.
Avoid contact with skin.
Wash thoroughly after handling.
Repeated contact may be harmful.
Do not contaminate water, streams or fish ponds.
Do not use on crops to be used for food.
This product is toxic to fish. Keep out of fish streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

EPA Reg. No. 572-43

© 1974 Waltham Chemical Works, Inc.

Waltham, Mass.

Waltham Chemical Works, Inc.
(St. Louis, Missouri 63160)

DIRECTIONS FOR USE

TRUBAN is a soil fungicide for control of damping off, root and stem rot diseases caused by Pythium and Phytophthora. It may be used at time of seeding and transplanting either as a soil mix or as a drench for control of these problem fungi.

TRUBAN suspensions in water should be constantly agitated to assure uniform application of material. TRUBAN suspensions need not be protected from light. Use at recommended rates. Over dosage may result in toxicity to sensitive plants. If plants are suspect of being sensitive, apply fungicide to a small number to insure safety. TRUBAN should not be mixed with other pesticides or nutrients.

Thoroughly rinse TRUBAN residues out of all equipment after use.

Plant Type	Remarks
Bedding Plants Ageratum Impatiens Alyssum Marigold Aster Pansy Balsam Petunia Begonia Portulaca Calendula Salvia Carnation Snapdragon Celosia Stokesia Coleus Verbena Dahlia Vinca Dianthus Zinnia Foxglove	BEDDING PLANTS (soil 2" to 3" deep) Mix 4-6 ozs TRUBAN 30% W.P. with 100 gallons of water and apply to 800 sq ft of bed area (1 gal per 8 sq ft) and use at time of seeding and transplanting. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re treat at 4 to 8 week intervals if additional applications are necessary.
Foliage Plants Aqlaonema Anthurium Chinese Evergreen Dieffenbachia Nephthytis Peperomia Pilea Pothos Philodendron Spp	BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6" deep) Mix 8-12 ozs of TRUBAN 30% W.P. with 100 gallons of water and apply to 400 sq ft of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re treat at 2-3 month intervals to maintain disease control.
	BED GROWN PLANTS Mix 3 to 10 ozs with 100 gals. of water per 400 sq ft. or apply in sufficient amount to saturate the soil mixture. Re treat at 4 to 12 week intervals if necessary.
Container & Bedgrown HERBACEOUS - Easter Lily Carnation - Geranium Chrysanthemum WOODY Arctosaphylus - Rhamnus Ceonothus - Rhododendron Poinsettia - Southern Pine	CONTAINER & BEDGROWN PLANTS Mix 3 to 10 ozs with 100 gals. of water per 400 sq ft. or apply in sufficient amount to saturate the soil mixture, such as 1/2 pint per 6" pot. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re treat at 4 to 12 week intervals if necessary.
DRY SOIL MIX for Plants Listed Above Mix 1 1/2 to 3 ozs TRUBAN 30% W.P. with each cu. yd. of soil mixture. Ingredients for the soil mix should be free of large lumps and dry enough to prevent agglomeration during mixing. (Improved uniformity of blending will be assured if the TRUBAN is pre mixed with about five times its volume of dry soil.) Re treat if needed as indicated under appropriate soil drench recommendations.	

9-27-76
 TRUBAN



BURN OR BURY EMPTY BAG - KEEP OUT OF SMOKE
NOTICE

Mallinckrodt makes no warranty, express or implied, concerning the fitness of this material for any particular purpose except those which are contained on the TRUBAN label.

