

BEST DOCUMENT AVAILABLE

APR 10 1982

1/2/83

KOBAN® 30

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: HARMFUL IF SWALLOWED. Inhaled, or absorbed through skin. Always wear protective clothing and respirator when mixing and applying pesticides. May cause irritation. Avoid breathing spray mist. Avoid contact with skin, eyes and on clothing. **IN CASE OF CONTACT,** immediately flush eyes or skin with plenty of water. **GET MEDICAL ATTENTION** if irritation persists. Wash thoroughly with soap and water after handling. Wash contaminated clothing before re-use. Do not contaminate feed or food. Keep children and pets out of treated area during application.

ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO FISH. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

FOR PROFESSIONAL USE ONLY

KOBAN 30 is intended for use on ornamental turf which is subject to Pythium blight, damping off or cottony blight. It may be used at time of seeding to control damping off, or on established turf to prevent or control Pythium blight (cottony blight).

KOBAN 30 does not afford broad spectrum fungus disease control for Rhizoctonia (brown patch), Sclerotinia (dollar spot), Glomerospora (copper spot), and similar turf diseases. A fungicide program to control these diseases should be followed if they co-exist with Pythium blight.

For control of Pythium blight (cottony blight, grease spot, damping off) on turf:

Night temperatures above 65°F (18.3°C.) with high humidity are favorable for Pythium development.

Add KOBAN 30 to the spray tank with the agitator in operation. Disperse 2 to 4 1/2 oz. of KOBAN 30 in 5 gal. of water for each 1000 sq. ft. of turf to be treated. Use 2 oz./1000 sq. ft. when disease incidence is expected to be light and 4 1/2 oz./1000 sq. ft. for more severe conditions.

Apply uniformly to the turf area with a SPRAY-HAWK® mobile spray boom or other suitable equipment.

Re-treat in 5-10 days, depending upon climatic conditions, to maintain disease control. KOBAN 30 should not be mixed with other pesticides or nutrients.

For control of damping-off in newly seeded areas: Apply 7 to 9 oz. of KOBAN 30 in 5 to 10 gal. of water/1000 sq. ft. immediately after seeding. Irrigate promptly for improved soil penetration of the fungicide and to initiate seed germination. Re-treat at 5 to 7 day intervals at the 2 to 4 1/2 oz. rate in order to maintain a preventive disease control program after germination.

Use FUNGO® 50, KROMAD®, DUOSAN® or CAD-MINATE® to control a broad spectrum of diseases other than Pythium.

Grass Seed Treatment: To prevent Pythium blight add 4 1/2 to 7 oz. of KOBAN 30 to the seed treaters for each 100 lb. of grass seed. Blend until the KOBAN 30 is thoroughly dispersed in the seed.

NOTE: For smaller seeds use the higher rate of KOBAN 30.

IMPORTANT: Bags containing treated seed should bear a tag with the following statement: This seed has been treated with 30% Ethazole at the rate of 4 1/2 to 7 oz./100 lb. seed. **DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES.**

STORAGE AND DISPOSAL

STORAGE: Not sensitive to light. Stable at normal temperatures and humidities. Store in a dry place, at temperature above 45°F (7.2°C), in a well-closed container.

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into applicator equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

WARRANTY

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given hereon. **NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.**

Printed in U.S.A. & C.