

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER! Corrosive causes eye damage and skin irritation. IF IN EYES GET MEDICAL ATTENTION IMMEDIATELY. Wash with large amounts of water for at least 15 minutes. If eye irritation continues, seek medical attention. If on skin, wash thoroughly with large amounts of water. If swallowed, drink plenty of water and call a physician or poison control center. If inhaled, move to fresh air. If breathing difficulty occurs, call a physician or poison control center.

Inhalation: Use caution to avoid inhaling spray mist. Avoid breathing spray mist or dust.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to lakes, streams, ponds, or contaminated water bodies. Do not contaminate surface waters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use on stone, asphalt, concrete.

DIRECTIONS FOR USE

It is a violation of Federal (USA) law to use this product in a manner inconsistent with its labeling.

FOR PROFESSIONAL USE ONLY

TRUBAN is a soil fungicide for control of damping-off, root and stem rot diseases caused by Pythium and Phytophthora. It may be used at time of seedling and transplanting as a drench for control of these problem fungi in greenhouses and outdoors.

TRUBAN suspensions in water should be applied to the soil before application of material. TRUBAN suspensions need not be protected from light. Use at recommended rates. Overdosage may result in toxicity to sensitive plants. Plants are suspect of being sensitive. Apply fungicide at a small number to insure safety. TRUBAN should not be mixed with other pesticides or nutrients.

Thoroughly rinse TRUBAN residues out of all equipment after use. Always use protective clothing and respiratory when mixing and applying peat.

STORAGE AND DISPOSAL

STORAGE Not sensitive to light. Stable at normal temperature and humidity.

PROPOSITIONS Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL Do not contaminate water, food, or feed by storage or disposal.

4326

ACCEPTED **1 QT.**
(0.95 Liter)

SEP 1 1981

See panel directions for use of this product.
 EPA Reg. No. 612-124

FUNGICIDE

25% EMULSIFIABLE CONCENTRATE

For control of damping off, root and stem rot diseases caused by Pythium and Phytophthora in ornamental and nursery crops.

5-Ethoxy-3-trichloro-methyl-1,2,4-thiadiazole* 25%
 * 75%

KEEP OUT OF REACH OF CHILDREN

DANGER!

CORROSIVE

CAUSES EYE DAMAGE AND SKIN IRRITATION

HARMFUL IF SWALLOWED

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes immediately with large amounts of water and call a physician or poison control center. If eye irritation continues, seek medical attention.

IF SWALLOWED: Do not induce vomiting. Give 1 glass of water and call a physician or poison control center. If unable to drink, have person obtain medical attention immediately.

IF INHALED: Move to fresh air.

IF ON SKIN: Wash thoroughly with soap and water. If irritation occurs, call a physician or poison control center.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 612-124

MALLINCKRODT, INC.

PLANT TYPE	REMARKS
Bedding Plants	BEDDING PLANTS (soil 2 to 3" deep) Mix 4 fl. oz. TRUBAN 25% E with 100 gal. of water and apply evenly over the soil. Do not water immediately after treatment. Re-apply immediately if rain falls. Do not water again until 24 hours have passed since the fungicidal drench has been applied. Re-treat at 4 week intervals if additional application is necessary.
Foliage Plants	BED AND BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6" deep) Mix 4 fl. oz. TRUBAN 25% E with 160 gal. of water and apply evenly over the soil. Do not water immediately after treatment. Re-apply immediately if rain falls. Do not water again until 24 hours have passed since the fungicidal drench has been applied. Re-treat at 3 month intervals if disease continues.
Container & Bedgreen	CONTAINER AND BEDGROWN PLANTS Mix 4 fl. oz. with 100 gal. of water per 400 sq. ft. or apply amount to saturate the soil mixture such as 1 pint per 6' immediately. Add additional water equal to at least half the volume of the fungicidal drench for improved penetration of the fungicide. Re-treat at 4 week intervals if necessary.

WARRANTY

Malinckrodt, Inc., warrants that the product will be effective against the disease problems listed on the label when used according to directions on the label. This warranty does not cover damage to the product resulting from misuse or abuse. Malinckrodt, Inc., shall not be liable for any damages resulting from the use of this product.

Editor

STATEMENTS**4326****1 QT.****(0.95 Liter)****ACCEPTED**

SEP 1 1981

6-Ethoxy-3-trichloro-methyl-1,2,4-thiadiazole
is the general trade name
of the active ingredient in
this product. This pesticide
is registered in the U.S.A.
EPA Reg. No. 572-44

FUNGICIDE**25% EMULSIFIABLE CONCENTRATE**

For control of damping off root and stem rot diseases caused by *Pythium* and *Phytophthora* in ornamental and nursery crops.

6-Ethoxy-3-trichloro-methyl-1,2,4-thiadiazole*	25%
.....	75%

KEEP OUT OF REACH OF CHILDREN**DANGER!****CORROSIVE**

CAUSES EYE DAMAGE AND SKIN IRRITATION
HARMFUL IF SWALLOWED

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes immediately with plenty of water for at least 15 minutes. **GET MEDICAL ATTENTION**

IF SWALLOWED: Induce vomiting immediately giving two glasses of water and stirring finger down throat. Never give anything by mouth to an unconscious person. **OBTAIN MEDICAL ATTENTION IMMEDIATELY**

IF INHALED: Remove to fresh air.

IF ON SKIN: Wash thoroughly with soap and water. **DO NOT SCRUB**. **GET MEDICAL ATTENTION**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 572-44 EPA Est. N. 226

MALLINCKRODT, INC.

PLANT TYPE	REMARKS
Bedding Plants	BEDDING PLANTS (soil 2 to 3 deep) Mix 4 fl. oz. with 100 gal. of water per 400 sq. ft. or apply sufficient amount to saturate the soil medium such as peat or topsoil. Broadcast evenly over the surface of the soil or in trench or strip evenly throughout the length of the bed. Allow to soak in for 1 hour to 1 day before planting.
Foliage Plants	BED AND BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6 deep) Mix 4 fl. oz. with 100 gal. of water per 400 sq. ft. or apply sufficient amount to saturate the soil medium such as peat or topsoil. Broadcast evenly over the surface of the soil or in trench or strip evenly throughout the length of the bed. Allow to soak in for 1 hour to 1 day before planting.
Container & Bedgrown	CONTAINER AND BEDGROWN PLANTS Mix 4 fl. oz. with 100 gal. of water per 400 sq. ft. or apply sufficient amount to saturate the soil medium such as peat or topsoil. Broadcast evenly over the surface of the soil or in trench or strip evenly throughout the length of the bed. Allow to soak in for 1 hour to 1 day before planting.

WARRANTY

Mallinckrodt, Inc., warrants that this product will remain in good condition and perform satisfactorily under normal conditions of handling, storage and use for a period of one year from the date of manufacture.

THIS WARRANTY IS EXCLUSIVE AND IN LIEU OF ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.