

REMARKS	
1.0 Soil Seedling benches etc.	BEDDING PLANTS (soil 2" to 3" deep) Mix 3.4 oz. TRUBAN 25% EC with 100 gal. of water and apply 1/800 sq. ft. of bed area of spray per 8 sq. ft. and use at time of seeding and transplanting. Irrigate immediately with additional water equal to at least half the volume of the fungicidal spray. For improved soil penetration of the fungicide. Re-treat at 4-6 week intervals if additional applications are necessary.
1.1 Soil bedding benches etc. 1.2 Soil benches etc.	BED AND BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6" deep) Mix 4.8 oz. of TRUBAN 25% EC with 100 gal. of water and apply to 400 sq. ft. of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicidal spray. For improved soil penetration of the fungicide. Re-treat at 2-3 month intervals to maintain disease control.
1.3 Soil potting benches etc.	CONTAINER AND BEDGROWN PLANTS Mix 4.8 oz. with 100 gal. of water per 400 sq. ft. coverage. If there is a tank or container, calculate the coverage required per pot. Irrigate immediately with additional water equal to at least half the volume of fungicidal spray. For improved soil penetration of the fungicide. Re-treat at 4-12 week intervals if necessary.

WARRANTY

Other labels do not state that the product contains Tridemorphine. This label does. It is recommended that you read all labels before using this product. If you have any questions concerning the use of this product, contact your distributor or manufacturer. The manufacturer will not be liable for damage resulting from misuse of this product. THE MANUFACTURER MAKES NO WARRANTY OR GUARANTEE AS TO THE MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

1.0 — SOLVENT N.O.S. (Petroleum Aromatic Hydrocarbons)
NA1993

SHAWMEN