

5001

2 LBS.



ACTIVE INGREDIENT:
 Dimethyl 4,4'- α -phenylenebis
 (3-thioallophanate)* 50%
 INERT INGREDIENTS: 50%
 Total 100%
 *Thiophanate M

CAUTION!

KEEP OUT OF REACH OF CHILDREN
 May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothes before reuse. Do not contaminate feed or food.
 Keep children and pets off treated area until material has been washed into the soil and area is completely dry.
 Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

ANTIDOTE

External: If spilled on skin, wash off immediately with soap and water. If in eyes, flush with plenty of water.
Internal: Induce vomiting with a mixture of salt, soap or mustard in warm water. CALL A PHYSICIAN IMMEDIATELY.
 Burn or bury empty containers, stay out of smoke; never reuse them.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MALLINCKRODT CHEMICAL WORKS

St. Louis, Missouri 63160

EPA Registration No. 372-45-AA

Aug 30, 1973
 372-45

BEST DOCUMENT AVAILABLE

43

GENERAL INFORMATION

FUNGO 50 is a wettable powder for use on golf course greens, tees and fairways of Bentgrass, Bluegrass, Perennial ryegrass, Fescue, Paspalum, St. Augustine grasses or their mixtures. FUNGO 50 is to be used for the prevention and control of the below mentioned diseases.

DIRECTIONS FOR USE

Add the recommended amount of FUNGO 50 to a partially filled spray tank under agitation. Complete filling the tank to the desired volume. Continuous agitation is necessary to insure even suspension of the wettable powder. Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a boom type sprayer or other power spray equipment. Apply after mowing or avoid mowing for twelve hours after application.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually two gallons per 1000 sq. ft. of turf area. When treating golf greens always treat aprons and approaches. Under conditions of severe disease the interval between applications may be shortened to 5-7 days.

For the following diseases the spray should dry on the leaves, with no "watering in":

DOLLAR SPOT (*Sclerotinia homocarpa*)—Apply FUNGO 50 at 1 oz. per 1000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals as long as needed. As a curative, where preventive schedule is not followed, use 2 ozs. per 1000 sq. ft.

GOULD SPOT (*Gloeocercospora sorghi*), **LARGE BROWN PATCH** (*Rhizoctonia solani*), **RED THIRTY** (*Corticium fuckelii*), **TRI-ANGULAR PATCH** (*Fusarium nigralis*)—Apply FUNGO 50 at 2 ozs. per 1000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals. Retreat at 5 to 7 day intervals under severe disease conditions.

For the diseases below, "watering in" is required:

USARIAL BLIGHT (*Fusarium roseum*)—Apply FUNGO 50 at 4-6 ozs. per 1000 sq. ft. Water into root zone by irrigation thoroughly immediately after application. Apply two applications at 10-14 day intervals, beginning application when the disease first appears.

SMUT (*Ustilago striiformis*)—Apply FUNGO 50 at 4-6 ozs. per 1000 sq. ft. Water into root zone thoroughly immediately after application. Apply in October or early spring just before grass begins growth. Do not graze treated areas or feed clippings to livestock.


07/73

BEST DOCUMENT AVAILABLE

44

372-45

5991



SY

GENERAL INFORMATION

FUNGO 50 is a wettable powder for use on golf course greens, tees and fairways of Bentgrasses, Bluegrasses, Perennialgrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. FUNGO 50 is to be used for the prevention and control of the below mentioned diseases.

DIRECTIONS FOR USE

Add the recommended amount of FUNGO 50 to a partially filled spray tank under agitation. Complete filling the tank to the desired volume. Continuous agitation is necessary to insure even suspension of the wettable powder. Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a boom type sprayer or other power spray equipment. Apply after mowing or avoid mowing for twelve hours after application.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually five gallons per 1000 sq. ft. of turf area. When treating golf greens always treat aprons and approaches. Under conditions of severe disease the interval between applications may be shortened to 5-7 days.

For the following diseases the spray should dry on the leaves, with no "watering in":

DOLLAR SPOT (*Sclerotinia homocarpa*) -- Apply FUNGO 50 at 1 oz. per 1000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals as long as needed. As a curative, where preventive schedule is not followed, use 2 ozs. per 1000 sq. ft.

COPPER SPOT (*Gloeocercospora sorghi*), LARGE BROWN PATCH (*Rhizoctonia solani*), RED THREAD (*Corticium fuciforme*), FUSARIUM PATCH (*Fusarium nivale*) -- Apply FUNGO 50 at 2 ozs. per 1000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals. Retreat at 5 to 7 day intervals under severe disease conditions.

For the diseases below, "watering in" is required:

FUSARIUM BILTCH (*Fusarium roseum*) -- Apply FUNGO 50 at 4-8 ozs. per 1000 sq. ft. Water into root zone by irrigating thoroughly immediately after application. Apply two applications at 10-14 day intervals, beginning applications when the disease first appears.

STRIPED SMUT (*Ustilago striiformis*) -- Apply FUNGO 50 at 4-8 ozs. per 1000 sq. ft. Water into the root zone thoroughly immediately after application. Apply in October or early spring just before grass begins growth. Do not use treated areas or feed clippings to livestock.

Cameraman's Note

Poor Copy