

ł

# KEEP OUT OF REACH OF CHILDREN WARNING!


### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Never give anything by mouth to an unconscious person. OBTAIN MEDICAL ATTENTION IMMEDIATELY.

IF INHALED, Induce coughing. IF ON SKIN, Wash off with soap and water.

IF IN EYES. Flush eyes with water.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.



11

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNINGI Causes eyas and skin Irritation. Do not get in eyes, on skin ar on clothing. Harmiul if swallowed or inhalod. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets off treated area until dry or material has washed into the soil. Always wear protective clothing and respirator when mixing and applying pesticides.

### ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO FISH. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Add recommended amount to a partially filled spray ank under agitation. Complete filling to the desired volume. Keep agitated to insure even suspension of material Make fresh daily. Spray uniformly over the area with a boom-type or other power sprayer. Apply after mowing or avoid mowing for 12 hours after application. Apply in sufficient water to obtain thorough coverage, usually 5 gai. per 1000 sq. ft. of turf area. Always treat aprons and approaches to golf greens. Under conditions of severe disease, apply every 5 to 7 days. Note to user: Do not apply to areas likely to be grazed by livestock and de not feed clippings to livestock or poultry.

**DUOSAN**\* is a weltable powder for use on golf course greens, tees, lawn and fairways of Bentgrasses, Bluegrasses, Fescues, Bermudagrasses, Ryegrasses, St Augustine grass or their mixtures to be used for the prevention and control of the following diseases:

> DOLLAR SPOT Scierolinia homesocarpa -- Apply at 3 oz per 1000 sq ft. beginning when the disease first appears and continue at 10-14 day intervals as needed. Retreat at 5-7 day intervals under severe conditions

BROWN PATCH Rhizocionia solani COPPER SPOT Glosocarcos-

pora sorghi

RED THREAD Corticium fuciforme

LEAF SPOTS Helminthosporium spp

FUSARIUM PATCH Fusarium nivale

RUST Puccinia ceronata

Apply at 4-6 oz. per 1000 sq ft. when disease appears and continue at 10-14 day intervals. Use 5-7 day intervals for severe disease.

## STORAGE AND DISPOSAL

STORAGE: Do not store at a temperature exceeding 122°F (50°C). Store in a dry place in a wellclosed container

PROVIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. PESTICIDE DISPOSAL: Rinse container three times with water. Drain each rinse into a sprayer or

mix tank. Dispose of all material and rinsate not used in accordance with governmental regulations

CONTAINER DISPOSAL: Discard empty containers in accordance with governmental regulations

WARRANTY

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices inature of soil or polying medium, the disease problem condition of the crop incompatibility with othercoemicals out specifically recommended and other influencing factors in the use of this product are beyond the control of the selles. Buyer emicrose all helps of the structure of helps of the selles of the se