

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before smoking or eating. Wash contaminated clothing before reuse. Do not contaminate feed or food. Do not permit children or pets to go into the treated areas until spray has completely dried.

ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO FISH and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICEPERSONS

DYMET® is useful against insect pests of turf, ornamentals and nursery crops. For use in greenhouses and outdoors. Do not apply to food crops grown in the greenhouse. Do not use in barns or dwellings.

On turf apply 1 1/2 to 3 fl. oz. in 5 gal. of water per 1,000 sq. ft. for control of bluegrass billbugs, southern chinchbug, fairy chinchbug, sod webworm and dingy cutworm. Apply 6 to 12 fl. oz. in 5 gal. of water per 1,000 sq. ft. for control of white grubs of southern masked chaffer and crane fly larvae. Irrigate to insure penetration of soil.

On ornamentals and nursery crops, apply 2 to 4 fl. oz. of water (1 to 1 1/2 fl. oz. in 1 gal.) for control of the following insects: aphids, caterpillars, mealy bugs, whiteflies, scale, leafhoppers, thrips, armyworms, webworms, leaf beetles, sawflies, lacebugs, whiteflies, earwigs, and grasshoppers. Spray thoroughly to runoff. Begin application when insects first appear and repeat at 7 to 14 day intervals as needed.

STORAGE AND DISPOSAL

STORAGE: Not sensitive to light. Store at 50° to 80° F. in cool, dry place. Store in original container. Do not use if container is damaged.

PROHIBITIONS: Do not use to fumigate areas. Do not use for storage or disposal. Do not use for disposal of waste.

PESTICIDE DISPOSAL: Pesticide should be disposed of in accordance with local, state or federal regulations. Do not dispose of in water that cannot be used for drinking or irrigation.

CONTAINER DISPOSAL: Containers should be disposed of in accordance with local, state or federal regulations. Do not reuse containers.

5731

Mallinckrodt

5 GAL.
(18.9 Liter)



* Equivalent to 17.6% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.4% other isomers and reaction products

** Diazinon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES. Flush eyes with plenty of water for 15 minutes. Get medical attention.

IF INHALED. Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN. Wash off with soap and water. Remove contaminated clothing and wash before reuse.

IF SWALLOWED. Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

Note to Physician. If signs of cholinesterase inhibition occur. Atropine is antidotal.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA EST. NO. 602-MO-1

EPA REG. NO. 572-54

MANUFACTURED FOR

MALLINCKRODT, INC.

ST. LOUIS, MISSOURI 63147

The following host plants are tolerant to DYMET in the prescribed application range (2 to 3 qt. per 100 gal. of water):

HERBACEOUS	WOODY
African Violet	Araucaria
Aglaonema	Arbovitae
Alternanthera	Ardisia
Aphelandra	Aucuba
Aralia	Azalea
Asparagus sprengeri	Birch
Begonia	Boxwood
Caladium	Camellia
Capsicum	Chamaecyparis
Cineraria	Chamaedorea
Cissus	Cherry
Coleus	Citrus
Cordatum	Cotoneaster
Creeping Charlie	Crape Myrtle
Croton	Douglas Fir
Dreftbachia	Elm
Dwarf Jade Plant	Euonymus
Episcia	Fatsia
Geranium	Ficus
Gladiolus	Fountain Grass
Gynura	Gardenia
Hemigraphis	Hazelnut
Homocladium	Ilex
Hoya Carnosa	Juniper
Impatiens	Ligustrum
Ivy	Lilac
Jade Plant (Crassula)	Liriodendron
Lily	Locust
Maranta	Magnolia
Marigold	Maple
Mondo Grass	Mimosa
Nepenthes	Nandina
Ochrosia	Ornamental palm
Peperomia	Osage
Peristrophe	Pecan
Petunia	Philadelphus
Philodendron	Plum
Pothos	Prunella
Salvia	Pine
Sansevieria	Pittosporum
Saxifraga	Pomegranate
Schumbergera	Prunus
Sedum	Pyracantha
Snapdragon	Rose
Spathiphyllum	Sago palm
Wandering Jew	Schefflera
	Sycamore
	Taxus
	Thuja
	Viburnum
	Weigelia
	Yucca

ACCEPTED
DEC 21 1983
Under the Federal Insect, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 372-54

Varieties and species of Calathea, Celosia, Chrysanthemum, Copper Plant, Dracaena, Gloxinia, and Poinsettia are sensitive to DYMET at higher rates.

WARRANTY