

- **WARNING! MAY BE POISONOUS IF ABSORBED THROUGH SKIN. HARMFUL IF SWALLOWED OR INHALED. CAUSES EYE IRRITATION.** Always wear protective clothing and respirator when mixing and applying pesticides. May cause irritation. Avoid breathing spray mist. Avoid contact with skin, eyes and on clothing. **IN CASE OF CONTACT**, immediately flush eyes or skin with plenty of water. **GET MEDICAL ATTENTION** if irritation persists. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets out of treated area during application.

THIS PRODUCT IS TOXIC TO FISH. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not use or store near heat or open flame

IT IS A VIOLATION OF FEDERAL (U.S.A.) LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

TRUBAN FLOWABLE is a soil fungicide for control of damping off, roo and stem rot diseases caused by *Pythium* and *Phytophthora*. It may be used at time of seeding and transplanting as a drench for control of these problem fungi in greenhouses and outdoors. **TRUBAN FLOWABLE** suspensions in water should be occasionally agitated to assure uniform application of material. **TRUBAN FLOWABLE** suspensions need not be protected from light. Use at recommended rates. Overdosage may result in injury to sensitive plants. If plants are suspect of being sensitive, apply fungicide to a small number for a safety test. **TRUBAN FLOWABLE** should not be mixed with other pesticides or nutrients.

Thoroughly rinse TRUBAN FLOWABLE residues out of all equipment after use. Always wear protective clothing and respirator when mixing and applying pesticides.

STORAGE AND DISPOSAL

STORAGE Store at 2-8°C (36-46°F) in original container. Do not freeze. Do not use if frozen.

PROHIBITIONS

PESTICIDE DISPOSAL _____

CONTAINER DISPOSAL : _____

**1 GAL.
(3.8 Liter)**

ACTIVE INGREDIENT
5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole* .. 40.7%

INERT INGREDIENTS 59.3%

**MAY BE POISONOUS IF ABSORBED THROUGH SKIN.
HARMFUL IF SWALLOWED OR INHALED
CAUSES EYE IRRITATION**

STATEMENT OF PRACTICE TREATMENT

" SWALLOWED DO NOT INDUCE VOMITING

GET MEDICAL ATTENTION

IF ON SKIN: Wash with plenty of water. Do not use solvents, oil-based products, or abrasives.

IF IN EYES

IF UNCONSCIOUS

CONTACT YOUR LOCAL POISON CONTROL CENTER HOSPITAL OR PHYSICIAN IMMEDIATELY
See Side Panels for Additional Precautionary Statements

Mallinckrodt, Inc.

EPA Reg No 3/2-59

EPA ESI No 1258-MS-1

PLANT TYPE		REMARKS
Bedding Plants		BEDDING PLANTS (soil 2" to 3" deep)
Agave Calceolaria Canna Chrysanthemum Gladiolus Dianthus Larkspur Marigold	Petunia Fuchsia Hibiscus Salvia Scabiosa Veronica Vinca Zinnia	Mix 2 1/2 lb of TRIBAN FLOWABLE with 100 gal of water and apply to 800 sq ft of bed area and use at time of seeding and transplanting. Irrigate immediately with additional watering sufficient to saturate the volume of the fungicide and direct the fungicide into penetration of the fungicide. Repeat at 4-8 week intervals if additional application is necessary.
Foliage Plants		BED AND BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS (peat and other high organic growth media 6" deep)
Agave Anthurium Begonia Cactus Caladium Calla Chamaelirium Chlorophytum Crassulaceae Daphne Dracaena Euphorbia	Ficus Maranta Nephrolepis Peperomia Philodendron Platanus Rhododendron Spathoglottis Zinnia	Mix 2 1/2 lb of TRIBAN FLOWABLE with 100 gal of water and apply to 800 sq ft of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicide. In the improved soil penetration of the fungicide. Repeat at 2-3 month intervals for maximum disease control.
Container & Bedgrown		CONTAINER AND BEDGROWN PLANTS
Agave Anthurium Begonia Cactus Caladium Calla Chamaelirium Chlorophytum Crassulaceae Daphne Dracaena Euphorbia	Ficus Maranta Nephrolepis Peperomia Philodendron Platanus Rhododendron Spathoglottis Zinnia	Mix 2 1/2 lb of TRIBAN FLOWABLE with 100 gal of water and apply to 800 sq ft of bench area. Irrigate immediately with additional water equal to at least half the volume of the fungicide. In the improved soil penetration of the fungicide. Repeat at 2-3 month intervals for maximum disease control.

WARRANTY

THE USER OF THIS PRODUCT ASSUMES ALL RESPONSIBILITY FOR THE PROPER USE OF THE PRODUCT. THE USER SHALL BE RESPONSIBLE FOR THE PROPER APPLICATION OF THE PRODUCT AND FOR THE PROTECTION OF THE ENVIRONMENT. THE USER SHALL BE RESPONSIBLE FOR THE PROTECTION OF THE ENVIRONMENT. THE USER SHALL BE RESPONSIBLE FOR THE PROTECTION OF THE ENVIRONMENT.