

DIRECTIONS FOR USE

General Information

Residex is a 100% active ingredient concentrate that must be diluted with water before use. For soil applications, Residex must be diluted with water.

Termite Control:

1. Prior to Slab Type Construction
 - a. Apply 1 gallon per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms, utility entrances, etc., have adjoining slab on foot foundations.
 - b. Apply 1 gallon per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms, utility entrances, etc., have adjoining slab on foot foundations.

2. Crawl Space Houses

- a. Apply 2 gallons per 10 square feet in a trenching operation to initial areas under the house, including the inside of foundation walls and around pipes and utility entrances. If trench is deeper than 15 inches, apply 4 gallons per 10 linear feet.
 - b. Apply 2 gallons per 10 square feet in a trench along the outside of the foundation wall, including entrances, platforms, and porches where the foundation is not more than 15 to 18 inches deep. If over 15 to 18 inches, but not over 30 inches deep, apply 4 gallons to 5 linear feet of trench. A 30 inch trench should be dug for buildings with deep foundations. Do not dig below the top of the footings.
- Apply 1 gallon per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms, utility entrances, etc., have adjoining slab on foot foundations.

Lawn and Turf Pests

For soil and lawn insects other than termites, Residex should be diluted in water and applied at the following rates:

- Ants: 1.5 pints of Residex 8 lb. per acre
- Chiggers: 1 pint of Residex 8 lb. per acre
- Earwigs: 1 pint to 1 gal. of Residex 8 lb. per acre
- Grubs: 1.5 pints of Residex 8 lb. per acre
- Japanese Beetles: 1 gal. of Residex 8 lb. per acre
- Mole Crickets: 1 gal. of Residex 8 lb. per acre
- White Grubs: 1 gal. of Residex 8 lb. per acre

For treatment of the above insects, sufficient water must be used to obtain coverage. It is strongly recommended that after application, the lawn areas be heavily watered so that the Chlordane emulsion washes into the ground.

MIXING INSTRUCTIONS FOR SOIL TREATMENT
Although water of any temperature may be used, Residex 8 lb. emulsion has more easily in warm water. If extremely cold water is used, the emulsion must be stirred until capable of obtaining a uniform stable emulsion for application.

residex

CHLORDANE 8 LB.

EMULSIFIABLE CONCENTRATE

Containing 8 lbs. Chlordane per gallon.

WARNING: KEEP OUT OF REACH OF CHILDREN
(See Side Panel for Additional Warning Statements)

ACTIVE INGREDIENTS:

*Technical Chlordane 74%

INERT INGREDIENTS: 26%

Total 100%

*Equivalent to 44.4% octachloro-4, 7-methanotetrahydroindane and 29.6% related compounds

FOR CONTROL OF SUBTERRANEAN TERMITES, ANTS, CHIGGERS, WHITE GRUBS, EARWIGS, CHINCH BUGS, SOD WEB WORMS, MOLE CRICKETS, JAPANESE BEETLE LARVAE AND NON-RESISTANT ROACHES.

E.P.A. Reg. No. 3/3 26

E.P.A. Est. No. 3/3 NJ 1

MANUFACTURED BY

residex

CORPORATION

225 TERMINAL AVENUE • CLARK, N.J. 07066

ACCEPTED
JUL 14 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for use pesticide registered under F.F.D. No. 149-010

For 1 gallon of dilute at the rate of 1 gal. of 3 lb. + 90 gallons of water = 100 gal. of 1 lb. emulsion. INSTRUCTIONS FOR COCKROACH CONTROL by Cracks and Crevice treatment.

Use a 2:1 emulsion. For 2 gallons of dilute at the rate of 2 gal. of Residex 8 lb. + 90 gallons of 100 gal. of 1 lb. emulsion.

DO NOT USE IN FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT. Do not use in food areas where food is being processed, prepared or served.

APPLICATIONS WITHIN FOOD AREA FOOD HANDLING ESTABLISHMENTS

LIMITED TO CRACK AND CREVICE TREATMENT ONLY. Apply a fine stream of emulsion directly into cracks and crevices such as expansion joints between different elements of construction, junction boxes or switch boxes, drains or hollow equipment legs where cockroaches hide. Care should be taken to avoid depositing product onto exposed surfaces or introducing material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

WARNING

This solution, if brought into contact with skin, cause toxic symptoms. Avoid breathing spray and skin contact. In case of spillage on the skin with soap and water. Avoid contamination of foodstuffs. Do not use on household pets or birds. Avoid getting into eyes. Harmful if swallowed.

The product is toxic to fish and wildlife. Birds and treated areas may be killed. Keep out of body of water. Apply this product only as shown on this label. Do not contaminate water by cleaning or disposal of wastes.

Do not apply where there is a risk to crops and where weather conditions favor drift to treated areas.

Keep children and pets off treated areas until material has washed into the soil and ground. Do not reuse empty drums. Return to manufacturer or destroy by perforating or crushing. Bury in a safe place away from water.

Antidote:

EXTERNAL: In case of spillage on skin promptly with soap and water.

INTERNAL: Emetic of 1 tablespoonful must be followed by 1 glass of water. Call a physician.

NET CONTENTS