

PRODUCT NUMBER
3355



THE RIGHT APPROACH TO GOOD TURF

CHIPCO® 26019 FUNGICIDE

CHIPCO 26019 FUNGICIDE



**A Foliar Applied Fungicide for the Prevention and Control
of Certain Diseases of Turfgrass**

ACTIVE INGREDIENT:

3-(3,5-dichlorophenyl)-N-(1-methylethyl)	
-2,4-dioxo-1-imidazolidinecarboxamide	50.0%
INERT INGREDIENTS	50.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 359-685
EPA Est. No.

**NET CONTENTS
2 POUNDS
.907 Kg.**

Manufactured by
RHÔNE-POULENC CHEMICAL COMPANY
AGROCHEMICAL DIVISION
RHÔNE-POULENC INC.
Monmouth Junction N.J. 08852

1111

1111

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

Disease	Interval of Applications	RATE
		Oz. Product 1000 ft ²
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	GREENS AND TEES: Repeat at 14-21 day intervals as long as required	1.5 - 2.0
Brown Patch (<i>Rhizoctonia solani</i>)	FAIRWAYS and OTHER TURF AREAS. Repeat at 22 - 28 day intervals as long as required.	NOTE: On Fairways, for Dollar Spot control, use 1 oz./1,000 Ft ² .
Helminthosporium: Leaf Spot and Melting Out (<i>Helminthosporium spp.</i>)		
Fusarium Blight (<i>Fusarium roseum</i>)	Use only preventative foliar applications when conditions first become favorable for disease development. Two additional applications should be made at 2 to 3 week intervals.	4.0
Fusarium Patch (<i>Fusarium nivale</i>) PACIFIC NORTHWEST ONLY (West of the Cascade Mountains)	Repeat at 14-21 day intervals as long as required.	2.0 - 4.0
Gray Snow Mold (<i>Typhula spp.</i>) Pink Snow Mold (<i>Fusarium nivale</i>)	One application before first permanent snow cover. If possible, another application during a midwinter thaw.	2.0 - 4.0

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24 hour waiting period following treatment is preferred. Do not mix with any sticker, extender, or wetting agent. Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock or poultry.

Local conditions may affect the use of this fungicide. Consult State Agricultural Extension or Experiment Station specialists for specific recommendations, consistent with label, for local pest problems.

This product conforms to the chemical description on the label thereof and is reasonably fit for purpose stated on such label only when used in accordance with directions under normal use conditions. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, handling of this material, not in strict accordance with directions given herewith.

ACCEPTED

NOV 18 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act
or amendments thereto, this pesticide
meets the requirements of
EPA Reg. 263.655