

PN 25 600-64
113

A SCOOP IS ATTACHED FOR MEASURING CONVENIENCE

B-NINE®

CONTROLS HEIGHT,
PROMOTES FLOWERING OF PLANTS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Do not allow spray solution to touch skin. Wear protective clothing to protect hands, arms and legs, but when diluting and applying this product. Avoid breathing dusts. Wear a protective mask during mixing/loading operations. Wash hands and face thoroughly with soap and water after use or before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands.

Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance, contact your Regional Office of the EPA.

Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

ACCEPTED
DEC 10 1984
Under the FIFRA Insecticide, Fungicide, and Herbicide Act as amended, for the pesticide registered under EPA Reg. No. 400-69



DIRECTIONS FOR USE

Each formulator is responsible for obtaining EPA registrations for its formulated products. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not reenter treated areas until 24 hours after application, unless wearing protective clothing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE — Store in a dry location. Use spray and stock solutions within 24 hours. Do not store solutions. Immediate use is appropriate, if another component is added to spray mixture.

PESTICIDE DISPOSAL — Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paper: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOT FOR REFORMULATION

COMPOSITION

Active Ingredient: (% by weight)	
Daminozide(butanedioic acid mono (2,2-dimethylhydrazide))*	85.0
Inert Ingredients:	15.0
TOTAL:	100.0

*U.S. Patents 3,240,799 - 3,334,991

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg white or gelatin solution; if not available drink large amounts of water. Avoid alcohol. Call a physician immediately.

IF ON EYES OR SKIN: Wash with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before use.

IF INHALED: and abnormal symptoms develop see a physician.

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

Mfg. by **UNIROYAL CHEMICAL** — Division of **UNIROYAL, Inc.**, Naugatuck, Connecticut 06770

EPA REG. NO. 400-69

EPA EST. NO.

B-NINE is a Registered Trademark of Uniroyal, Inc.

©Copyright 1983, Uniroyal, Inc.

9-19-84

Lot. No.

NET CONTENTS:

APPLICATION NOTES

Do not add additional wetting agent or mix with insecticides or fungicides as burning of leaves may result.

Best results are obtained if plants are watered before spraying. Foliage should be dry when spray is applied. Wilted plants do not readily absorb B-NINE and poor results might be obtained if wilted plants are sprayed. Do not syringe for 18 to 24 hours after spraying to allow the chemical to enter the plant.

To measure B-NINE, use the large end of the scoop provided.

B-NINE® DILUTION TABLE

% B-NINE DESIRED	LEVEL SCOOPS per GALLONS** of solution		Wgt. Measure for one (1) gallon of solution	
	Scoops	Gallons	Ounces (overdrams)	Grams
0.15	1	3/4	0.24	6.7
0.25	2	1	0.40	11.1
0.37	3	1	0.60	16.7
0.50	4	1	0.80	22.3
0.75	6	1	1.20	33.4
1.00	8	1	1.60	44.5

Approximate dosage only. Use weight measurements wherever possible.

NOTE: Stock solutions prepared from B-NINE should not be held for more than 24 hours.

Measure the amount of B-NINE shown in an empty one-gallon container. Add enough warm water with stirring to make one gallon of solution. As B-NINE is soluble in water, no additional mixing is required.

GENERAL INFORMATION

B-NINE is dissolved in warm water then sprayed on the plant foliage using standard spray equipment. B-NINE passes into the leaf, then moves within the plant to reduce stem elongation. Treated plants develop a shorter, more compact habit of growth when compared to untreated plants. (See dilution table for converting percent concentration to dosage of B-NINE per gallon.)

DIRECTIONS FOR USE

Pet Chrysanthemums

To produce compact potted, branched (pinched) plants, spray to runoff 2 weeks after pinching with a 0.25% solution of B-NINE

If additional retardation is desired, a second application of 0.25% B-NINE may be applied 3 weeks after the first application. It is suggested that a "MEDIUM" schedule calling for shading and pinching on the same day be used for "TALL" varieties. This will eliminate the open, relatively leafless appearance often associated with the use of delayed pinch schedules.

Cut Chrysanthemums

To retard neck stretching just below the flower on potted and cut flowers, spray the top 1/3 of the foliage to runoff with 0.25% B-NINE two days after disbud-ding.

Azaleas — Greenhouse Plants

To promote development of additional buds on plants grown for forcing, spray when new growth following final shaping is 1-2 inches with either:

1. Two applications of 0.15% B-NINE at one-week intervals.
2. One application of 0.25% solution of B-NINE

Azaleas — Nursery Plants

To produce compact plants for landscape use, spray the foliage to runoff with a 0.37% solution of B-NINE during the fourth week in July. The final shaping of the plants should be done by early July to permit adequate regrowth prior to spraying.

Hydrangeas — Summer Growth

For those greenhouse varieties that produce excessive growth during the summer following pinching, spray to runoff with a 0.75% solution of B-NINE when the regrowths are 1-2 inches in length, usually in late July, but not after the first week in August.

Hydrangeas — Spring Growth

To retard spring forcing growth, spray with a 0.5% solution of B-NINE when the new growth has just started to unfold and four to five pairs of leaves are visible. Apply 2-4 weeks after the start of forcing but not later than this time.

Bedding Plants

Excessive stem elongation of petunias, marigolds, zinnias, asters, cosmos and salvia is prevented by a B-NINE spray made 2-3 weeks after transplanting, or when the plants begin to elongate (stretch). Use a 0.5% concentration sprayed to the drip point.

TO REDUCE LEAF INJURY OF PETUNIAS CAUSED BY AIR POLLUTANT, OZONE AND SULFUR DIOXIDE make a second application of 0.5% B-NINE one week after the first. B-NINE will not protect against leaf damage caused by the air pollutant PAN. Growth retardation will be more pronounced and there will be a delay in flowering from the second B-NINE application. Plants sprayed for pollution protection will be no shorter than those sprayed for compact growth.

POINSETTIAS

B-NINE reduces the elongation (stretching of poinsettias).
The following table is for pinched plants:

Pinch Date	Aug 15	Sept 1	Sept 15	
B-NINE Concentration	0.75%	0.5%	0.75%	
No. of applications	2	2	1	
The first application should be made when the new growth is 3-4 inches long. The second application, if suggested, should be made 2 weeks after the first application.				
The following table is for single stem plants				
Propagation date	Aug 1	Aug 15	Sept 1	Sept 15
First spray date	Sept 1	Sept 15	Oct 1	Oct 15
Concentration	0.75%	0.75%	0.5%	0.75%
Second spray date	Sept 21	Oct 1	Oct 15	None
Concentration	0.75%	0.5%	0.5%	—

The first application dates suggested are about one to two weeks after rooting. It is assumed that three weeks are required for rooting. Local conditions may require changing these suggested dates. But in all cases the first application should be made one to two weeks after rooting.

Gardenias

To produce compact plants for spring bloom, spray with 0.5% B-NINE when plants are about 2/3 of final marketing size. This is from mid-August to mid-September for the Florida area and during the first two weeks of December for other areas.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.